

Identification using Revealed Preferences in Linearly Separable Models*

Nikhil Agarwal, Pearl Z. Li and Paulo Somaini[†]

September 6, 2024

Abstract

Revealed preference arguments are commonly used when identifying models of both single-agent decisions and non-cooperative games. We develop general identification results for a large class of models that have a linearly separable payoff structure. Our model allows for both discrete and continuous choice sets. It incorporates widely studied models such as discrete and hedonic choice models, auctions, school choice mechanisms, oligopoly pricing and trading games. We characterize the identified set and show that point identification can be achieved either if the choice set is sufficiently rich or if a variable that shifts preferences is available. Our identification results also suggests an estimation approach. Finally, we implement this approach to estimate values in a combinatorial procurement auction for school lunches in Chile.

*We are grateful to Gabriel Weintraub for sharing the data used in this project. We thank Brent Hickman, Francesca Molinari, Parag Pathak, Anna Russo, and Whitney Newey for helpful comments. Yucheng Shang provided excellent research assistance. Agarwal acknowledges support from the Alfred P. Sloan Foundation (FG-2019-11484).

[†]Agarwal: Department of Economics, MIT and NBER, email: agarwaln@mit.edu. Pearl Z. Li: Department of Economics, Stanford University. Somaini: Stanford Graduate School of Business and NBER, email: soma@stanford.edu.

1 Introduction

Revealed preference arguments are central to estimating the distribution of agents’ payoffs. These arguments yield restrictions on preferences or payoffs based on an observed action (or choice). The typical argument assumes optimal behavior to derive implications of the requirement that any alternative action that an agent could have taken must yield a lower payoff. The consequences of the chosen and alternative action – in terms of realized allocations – are then used to identify or bound the payoffs from various allocations.

Such arguments have been applied to study a seemingly disparate set of models. The most immediate application is to consumer choice (e.g., [McFadden, 1973](#); [Rosen, 1974](#)).¹ However, arguments that are similar in spirit have been applied for other single-agent choice settings and a class of non-cooperative games. Single-agent examples include hedonic demand models (e.g., [Rosen, 1974](#); [Bajari and Benkard, 2005](#)) and monopoly pricing. Examples of non-cooperative games include games with incomplete information such as auctions ([Guerre et al., 2000](#)), school choice ([Agarwal and Somaini, 2018](#)), and trading or bargaining games ([Larsen and Zhang, 2018](#)); and games with full information such as oligopoly price setting where the objective is to identify marginal costs ([Berry et al., 1995](#); [Berry and Haile, 2014](#)). A common question is whether (the distribution of) agents’ payoff types is identified from the available data.

Our starting point is the observation that several models in the literature share a common structure. An agent can take an action $a \in \mathcal{A}$. The consequence of the action is described by an expected outcome $x \in \mathcal{X} \subseteq \mathbb{R}^J$ and an expected transfer $t \in \mathbb{R}$. Payoffs take a linear form and the agent maximizes

$$V(a; v) = v \cdot x_{\mathcal{A}}(a) - t_{\mathcal{A}}(a),$$

where $v \in \mathbb{R}^J$ is the agent’s preference type, and $x_{\mathcal{A}}(\cdot)$ and $t_{\mathcal{A}}(\cdot)$ are functions that map actions to outcomes and transfers respectively. We assume that the analyst knows (or can identify) $x_{\mathcal{A}}(a)$ and $t_{\mathcal{A}}(a)$ and observes each agent’s choices. An important requirement for conducting counterfactual analysis is that the cumulative distribution function (CDF) of the random variable v given observables z , denoted with $F_{V|Z}(v; z)$, is identified. In addition to this distribution, some cases also yield identification of the payoff type for each agent in the

¹We abstract away from the endogeneity of prices which is the focus of an important literature on identifying market demand ([Berry et al., 1995](#); [Berry and Haile, 2014](#)). We focus on cases where endogeneity can be perfectly controlled for by other means or where consumer-level price variation that is orthogonal to preferences is available (e.g. [Tebaldi et al., 2019](#)).

market.

This paper studies the identification of models with this linearly separable structure. Our framework and results allow for both discrete and continuous choice sets \mathcal{X} or a mixture of the two, and can incorporate both single-agent decision settings as well as a large class of non-cooperative games with independent private information. As we formally demonstrate in section 3, the hedonic demand model of Rosen (1974) and Bajari and Benkard (2005) maps to the model above if a is the good chosen, $x_{\mathcal{A}}(a)$ denotes the characteristics of the good, v denotes the vector of consumer preferences for the characteristics and $t_{\mathcal{A}}(a)$ denotes the pricing function. In the class of incomplete information non-cooperative games with independent and private types, $x_{\mathcal{A}}(a)$ and $t_{\mathcal{A}}(a)$ are expected outcomes and transfers, integrating over the strategies of other agents. For example, in the first-price auction studied in Guerre et al. (2000), a denotes the bid, $x_{\mathcal{A}}(a)$ denotes the probability of winning and $t_{\mathcal{A}}(a)$ denotes the expected payment. Similarly, in the model of Lewbel and Tang (2015), $a \in \{0, 1\}$ denotes a binary, $x_{\mathcal{A}}$ is the identity function, and $t_{\mathcal{A}}(a) = a \times z$ where z is a special regressor.

The main results characterize the identified set of distributions $F_{V|Z}(v|z)$ and derives conditions under which it is point identified. Point identification can be achieved in two cases. The base case is if the choice set is \mathcal{X} is sufficiently “rich” so that each choice is optimal only for a unique payoff type. This result implies identification in the hedonic price model as well as the first-price independent private value auctions. It also implies identification of marginal costs in oligopoly models if the demand function is known.

In the complementary case, where each choice is optimal for a set of payoff types, we show conditions under which variation in the observable z can be used to “trace out” the distribution of v . In this case, we require that z acts as a payoff shifter as in Lewbel (2000), although we can allow for a more general non-linear form by applying arguments similar to those in Allen and Rehbeck (2017). Our results imply identification in discrete choice models, bargaining models with discrete offers (Larsen and Zhang, 2018), and school choice models with or without strategic manipulation (Agarwal and Somaini, 2018).

These results therefore unify the analysis of identification in a large class of models, which have thus far been obtained using arguments customized for each model. While we do not aim to extend the analysis of identification in these models, we hope that the relatively sparse structure required for our results will be useful for new models. For instance, our model allows for a combination of discrete and continuous choices, which might be useful in some contexts – e.g., Aspelund and Russo (2023), which analyzes a scoring auction with multi-dimensional bids that include discrete and continuous components. We hope that our

general results and this shared structure will yield more immediate results for other models where identification results do not currently exist.

Our characterization of the identified set also suggests an estimation approach that is portable across contexts. In a first step, an analyst can use the revealed preference restrictions that we derive to bound the payoff type of each agent in the dataset based on the agent’s action. These restrictions yield a convex set to which the agent’s payoff type belongs. The second step then estimates the distribution of payoffs using an estimator of choice and further restrictions (if any).

As an illustrative application, we empirically analyze the procurement for public school lunches in Chile that is based on a combinatorial auction. This auction was analyzed in [Kim et al. \(2014\)](#), henceforth KOW. Our study uses the same dataset. It is well-known that the extreme high dimensionality of the choice set in a combinatorial auction presents several technical challenges. We show how our reformulation of the problem suggests an alternative solution to this dimensionality problem than the one taken in KOW.

Related Literature

A large literature that is not easily summarized applies revealed preference arguments to show identification of various models. We point the reader to several surveys for identification results pertaining to these models; for example, see [Berry and Haile \(2016\)](#) for demand models; [Athey and Haile \(2007\)](#) for auction models; and [Agarwal and Somaini \(2020\)](#) for school choice.

Our paper shares its focus on general revealed preference arguments with [Pakes \(2010\)](#), which also starts with revealed preference inequalities. The models and approaches are non-nested – [Pakes \(2010\)](#) allows for expectational errors in agents’ beliefs but places stronger functional form restrictions on the estimand, which is the expectation of v and z . We study the identification of the conditional distribution given by $F_{V|Z}$. In this sense, our estimand is similar to that in [Ciliberto and Tamer \(2009\)](#), although we pursue a non-parametric approach and consider conditions for obtaining point identification.

The illustrative application that we study is related to a small but growing literature on the analysis of combinatorial auctions. Papers that use bid data to estimate complementarities or substitutabilities between multiple objects that are auctioned off simultaneously include [Cantillon and Pesendorfer \(2007\)](#), [Gentry et al. \(2014\)](#) and [Xiao and Yuan \(2020\)](#).

Overview

Section 2 describes the notation of our model for single-agent problems and non-cooperative games, and maps the leading examples to our notation. Section 4 presents the main results

on identification and applies it to the examples. Section 5 presents the application to combinatorial auctions. Section 6 concludes. Proofs and lemmas not contained in the main text are in the Appendix.

2 Model

2.1 Notation and Agent Decisions

Consider an agent indexed by i who picks an action a from a set \mathcal{A} .² Let $x_{\mathcal{A}} : \mathcal{A} \rightarrow \mathbb{R}^J$ and $t_{\mathcal{A}} : \mathcal{A} \rightarrow \mathbb{R}$ be the functions mapping actions to outcomes and transfers respectively. Each action results in an **outcome** described by $x \in \mathcal{X} = \{x \in \mathbb{R}^J : x = x_{\mathcal{A}}(a) \text{ } a \in \mathcal{A}\}$ and an (expected) **payment** $t \in \mathbb{R}$. We do not restrict the cardinality of \mathcal{X} or of \mathcal{A} . By assuming that all agents face the same choice sets, our analysis effectively conditions on the value of $(x_{\mathcal{A}}, t_{\mathcal{A}})$ faced by an agent.

Agent i 's preference type is denoted $v_i \in \mathbb{R}^J$. Her (expected) utility from choosing $a \in \mathcal{A}$ is given by the linear form

$$v_i \cdot x_{\mathcal{A}}(a) - t_{\mathcal{A}}(a).$$

We assume that each agent chooses $a \in \mathcal{A}$ to maximize this indirect utility.³

Optimality implies that an agent with preference type v picks action a only if for all $a' \in \mathcal{A}$,

$$v \cdot x_{\mathcal{A}}(a) - t_{\mathcal{A}}(a) \geq v \cdot x_{\mathcal{A}}(a') - t_{\mathcal{A}}(a').$$

Thus, if a and a' are such that $x_{\mathcal{A}}(a) = x_{\mathcal{A}}(a')$ and $t_{\mathcal{A}}(a) < t_{\mathcal{A}}(a')$, then no agent picks a' . Define $t_{\mathcal{X}}(x)$ to be cost of choosing the outcome $x \in \mathcal{X}$:

$$t_{\mathcal{X}}(x) = \inf \{t_{\mathcal{A}}(a), a \in \mathcal{A} : x_{\mathcal{A}}(a) = x\},$$

where the infimum of an empty set is defined to be infinity. Observe that is it dominated to choose an action $a \in \mathcal{A}$ if $(x_{\mathcal{A}}(a), t_{\mathcal{A}}(a))$ is not in the graph of $t_{\mathcal{X}}(x)$.

We introduce two definitions that will be useful in the analysis. First, define the **convex hull** $t(\cdot) = \text{conv } t_{\mathcal{X}}(\cdot)$ of the function $t_{\mathcal{X}}(\cdot)$ to be the greatest convex function majorized by

²Agents are allowed to randomize between actions in the set \mathcal{A} . However, eventually the randomization results in a single action $a \in \mathcal{A}$.

³This formulation embeds scale and location normalizations because the expected utility is equal to $-t_{\mathcal{A}}(a)$ if $x_{\mathcal{A}}(a) = 0$.

the function (Rockafellar, 1970, page 36):

$$t(x) \equiv \inf \{t \mid (x, t) \in \text{conv epi } t_{\mathcal{X}}(\cdot)\},$$

where $\text{conv epi } t(\cdot)$ is the convex hull of the epi-graph of $t(\cdot)$.⁴ By definition, the convex hull of a function on \mathcal{X} is a convex function with domain $\bar{\mathcal{X}}$ given by the convex hull of \mathcal{X} .

Second, define the **subdifferential** $\partial t(x)$ of a convex function $t : \bar{\mathcal{X}} \rightarrow \mathbb{R}$ evaluated at x to be the set of all **subgradients** $v \in \mathbb{R}^J$ such that for all $x' \in \bar{\mathcal{X}}$, $t(x') \geq t(x) + v \cdot (x' - x)$. The subdifferential of a convex function is a non-empty convex set at every point in the interior of its domain. Further, if t is differentiable at x , then the subdifferential is a singleton containing only the gradient of t evaluated at x , $\nabla t(x)$.

Figure 1 illustrates the function $t_{\mathcal{X}}(\cdot)$, its convex hull $t(\cdot)$, and subgradients of $t(\cdot)$ at certain points. The horizontal axis denotes the potentially high-dimensional outcome space \mathcal{X} . The solid curve represents $t_{\mathcal{X}}(\cdot)$ and the dashed curve represents the parts where $t(\cdot)$ differs from $t_{\mathcal{X}}(\cdot)$. Specific pairs of x and t are labeled A through F , with co-ordinates $(x_A, t_A), \dots, (x_F, t_F)$. The graph of $t_{\mathcal{X}}(\cdot)$ contains all the points achievable by some action in \mathcal{A} . Thus, this graph contains A, B, C, E and F , but not the hollow points. The graph of $t(\cdot)$ does not contain the dominated point B but contains D and all points in the segment $D - E$ which can be achieved, in expectation, by randomizing between C and E , but not by choosing any particular action in \mathcal{A} . The function $t(\cdot)$ can exhibit upward discontinuities. As x approaches x_F , $t(x)$ approaches the hollow point below F . However, at x_F , $t(x_F)$ jumps upwards to point F .


Since $t_{\mathcal{X}}(\cdot)$ and $t(\cdot)$ are differentiable at x_A , the only element of the subdifferential is the slope of the dotted line through A . In contrast, these functions are not differentiable at C . The subdifferentials of $t(\cdot)$ consists of the slopes of all lines that are everywhere below the dashed and solid lines. The only element of the subdifferential of $t(\cdot)$ of D and E is the slope of the line through the points $C - E$.

2.2 Non-Cooperative Games

Although we began with the single-agent case, our framework also allows analysis of games that satisfy the following structure on payoffs and information. Let a_{-i} and v_{-i} respectively denote the actions and values of agents other than i . Let $\tilde{x}_i(a_i, a_{-i})$ be the outcome and $\tilde{t}_i(a_i, a_{-i})$ be the transfer for agent i as a function of the action profile (a_i, a_{-i}) . If the outcome

⁴The epi-graph of a function f is the set of points (x, y) such that $y \geq f(x)$.

Figure 1: Outcomes, Convex Hulls and Subgradients



Notes: The solid curve is the graph of $t_X(x)$ and the dashed curve is the graph of $t(x)$ for points where $t(x) < t_X(x)$. The points labeled A through F represent outcomes. When an outcome is represented by a solid circle, it belongs to the graph of $t_X(x)$ and results from an action $a \in \mathcal{A}$. An outcome represented by a hollow circle does not belong to the graph of $t_X(x)$. Such points do not yield from an action $a \in \mathcal{A}$.

is random conditional on the action profile, then interpret $\tilde{x}_i(a_i, a_{-i})$ and $\tilde{t}_i(a_i, a_{-i})$ be the corresponding expected values.

Assume that agent i 's payoff playing a_i when the other agents play a_{-i} is given by

$$v_i \cdot \tilde{x}_i(a_i, a_{-i}) - \tilde{t}_i(a_i, a_{-i}).$$

Let \mathcal{J}_i denote agent i 's information set, with $v_i \in \mathcal{J}_i$. The expected utility from playing a_i is given by

$$v_i \cdot \mathbb{E}[\tilde{x}_i(a_i, a_{-i}) | a_i; \mathcal{J}_i] - \mathbb{E}[\tilde{t}_i(a_i, a_{-i}) | a_i; \mathcal{J}_i],$$

where expectations are taken with respect to the distribution of actions a_{-i} of the other players given the information set \mathcal{J}_i . Uncertainty in this model can arise either because i expects its opponents to play a mixed strategy, because the types v_{-i} are private information from the perspective of agent i , or both.

This model fits our framework if each agent is best responding to the distribution of actions

played by others. Specifically, set

$$\begin{aligned} x_{\mathcal{A}}(a_i; \mathcal{J}_i) &= \mathbb{E}[\tilde{x}_i(a_i, a_{-i}) | a_i; \mathcal{J}_i] \\ t_{\mathcal{A}}(a_i; \mathcal{J}_i) &= \mathbb{E}[\tilde{t}_i(a_i, a_{-i}) | a_i; \mathcal{J}_i]. \end{aligned}$$

When analyzing games, we will assume that the two functions above are known to the analyst. A sufficient condition is that the outcome and the distribution of actions a_{-i} that agent i expects, $a_{-i} | \mathcal{J}_i$, is known or can be identified.

For example, consider the case in which agents' types are private information, each drawn independently from a distribution with CDF F_V . In this case, if the strategy profile $\sigma^*(v)$ constitutes a Bayesian Nash Equilibrium, then

$$x_{\mathcal{A}}(a_i) = \int \tilde{x}_i(a_i, \sigma_{-i}^*(v_{-i})) dF_{V_{-i}} \quad (1)$$

$$t_{\mathcal{A}}(a_i) = \int \tilde{t}_i(a_i, \sigma_{-i}^*(v_{-i})) dF_{V_{-i}}, \quad (2)$$

where the conditioning on the information set is dropped because it is irrelevant. Since play is described by a Bayesian Nash Equilibrium, agents have correct beliefs about the distribution of a_{-i} . The distribution of a_{-i} that each agent expects is identified from observation of the actions of all agents in independent and identically distributed instances of the game.

2.3 Regularity

We will make the following assumption to ensure that the optimization problem faced by each agent and their resulting choices are well-behaved:

Assumption 1. (i) The function $t_{\mathcal{X}} : \mathcal{X} \rightarrow \mathbb{R}$ is lower semi-continuous.

(ii) The set \mathcal{X} is non-empty and compact.

(iii) The random variable $X = X^*(V, \omega) \in \mathcal{X}$, where $X^*(\cdot, \cdot)$ is measurable with respect to V and $\omega \in \Omega$ and for all $\omega \in \Omega$

$$X^*(v, \omega) \in \arg \max_{x \in \mathcal{X}} v \cdot x - t_{\mathcal{X}}(x).$$

Parts (i) and (ii) above imply, by the extreme value theorem, that the solution to the problem $\max_{x \in \mathcal{X}} v \cdot x - t_{\mathcal{X}}(x)$ for each $v \in \mathbb{R}^J$ is attained for some element of \mathcal{X} . Part (i) rules out upward discontinuities in $t_{\mathcal{X}}(x)$ as in the hollow point below F in Figure 1 and part (ii)

rules out the hollow point D . Compactness of \mathcal{X} is either commonly satisfied or a weak restriction in the models discussed in section 3 below. For instance, in first-price auctions, \mathcal{X} is the unit interval; in the hedonic demand models, it restricts the characteristic space to be compact. Part (iii) assumes optimal behavior as well as a weak technical condition regarding the measurability of the map from types to outcomes. We allow for a randomization device ω for cases where there are multiple elements in \mathcal{X} that attain the maximum. We do not impose any restrictions on the joint distribution of ω across agents.⁵ Specifically, when $\arg \max_{x \in \mathcal{X}} v \cdot x - t_{\mathcal{X}}(x)$ is a set, then the agent is indifferent between multiple outcomes $x \in \mathcal{X}$ and part (iii) requires that the random variable X is generated from a measurable selector.

Under Assumption 1, $t(\cdot)$ is a lower semi-continuous proper convex function supported over a non-empty convex and compact set $\bar{\mathcal{X}}$. The conjugate function $t^*(v) = \max_{x \in \mathcal{X}} v \cdot x - t_{\mathcal{X}}(x)$ is also a lower semi-continuous proper convex function because the conjugate is the pointwise supremum over a family of affine functions with domain \mathbb{R}^J . Let $\mathcal{V}^\dagger \subseteq \mathbb{R}^J$ be the open set of points where $t^*(v)$ is differentiable. The set \mathcal{V}^\dagger is dense in \mathbb{R}^J and its complement is a set of measure zero Rockafellar (1970, Theorem 25.5). We define $\mathcal{X}^\dagger \subseteq \bar{\mathcal{X}}$ as the collection of gradients of t^* over \mathcal{V}^\dagger . More formally, $\mathcal{X}^\dagger = \{x \in \bar{\mathcal{X}} : x \in \partial t^*(v), v \in \mathcal{V}^\dagger\}$. If $v \in \mathcal{V}^\dagger$, then $x \in \partial t^*(v)$ is the unique solution to the problem $\max_{x \in \bar{\mathcal{X}}} v'x - t(x)$. Moreover, if $x \in \mathcal{X}^\dagger$, then there exist $v \in \mathcal{V}^\dagger$ such that $vx - t(x) > vx' - t(x')$ for all $x' \neq x$. Thus, \mathcal{V}^\dagger is the set of value with a strictly preferred outcome and \mathcal{X}^\dagger is the set of outcomes that are strictly preferred by some $v \in \mathcal{V}^\dagger$.

2.4 Observables

We follow the convention that upper-case letters denote random variables while lower case letters indicate specific values of the corresponding random variable.

Consider a dataset in which the analyst has access to a large sample of observations indexed by i from conditionally independent choices given a set of covariates Z . The analyst observes Z and the chosen outcome $X = X^*(V)$.

In addition, we assume that the analyst knows or observes the feasible outcomes \mathcal{X} and the function $t_{\mathcal{X}}(\cdot)$. Therefore, we assume that \mathcal{X} and $t_{\mathcal{X}}(\cdot)$ are either identified from the data or known from institutional details.

⁵Theorem 18.19 in Aliprantis and Border (2006) implies that a measurable selector satisfying part (iii) exists if \mathcal{X} is compact and $t_{\mathcal{X}}(\cdot)$ is continuous. This is the case because the maximand is a Caratheodory function: it is the difference between a bilinear function and a continuous function $t_{\mathcal{X}}(\cdot)$ of the argument x .

As discussed above, these outcomes and payoffs are generated by choosing actions $a \in \mathcal{A}$ with corresponding outcomes and transfers. Therefore, an alternative to observing X_i and having knowledge of $t_{\mathcal{X}}(\cdot)$ is that the actions A_i are observed and the functions $x_{\mathcal{A}}(\cdot)$ and $t_{\mathcal{A}}(\cdot)$ are known. The rest of the analysis treats X and $t_{\mathcal{X}}(\cdot)$ as observable although all of our conclusions carry over to the alternative case by setting $X = x_{\mathcal{A}}(A)$. In sum, we consider that the distribution of X conditional on Z is identified directly from the observed data.

We can also allow for \mathcal{X} and the function $t_{\mathcal{X}}(x)$ to vary as long as the sample includes many observations for a given pair $(\mathcal{X}, t_{\mathcal{X}}(\cdot))$. We omit variation in this object for notational simplicity because our analysis will be conditional on a fixed value.

In what follows, the conditioning on specific values of Z will be dropped from the notation except when we explicitly put together choices of agents with different values of the observables.

2.5 Identification

We follow the standard definitions of identification and falsifiability in the literature (e.g., [Athey and Haile, 2002](#); [Matzkin, 2007](#)). That is, a model is identified if the joint distribution of the model's primitives is uniquely determined by the joint distribution of observables. In our case, a model (\mathbb{F}, \mathbb{S}) is defined by a collection \mathbb{F} of joint distributions of V and Z , $F_{V,Z}$, and a collection \mathbb{S} of maps $\phi : \mathbb{F} \rightarrow \mathbb{H}$, where \mathbb{H} is the set of all joint distributions of X and Z , $F_{X,Z}$. We assume that the model is correctly specified. That is, the true value of $(F_{V,Z}, \phi)$ that generates the data belongs to (\mathbb{F}, \mathbb{S}) .

Our analysis will focus on identifying various features of the model, such as the joint distribution $F_{V,Z}$:

Definition 1. A feature $\psi : (\mathbb{F}, \mathbb{S}) \rightarrow \Omega$ is **identified** given the model (\mathbb{F}, \mathbb{S}) if and only if for any two pairs $(F_{V,Z}, \phi)$ and $(\tilde{F}_{V,Z}, \tilde{\phi})$ in (\mathbb{F}, \mathbb{S}) , $\phi(F_{V,Z}) = \tilde{\phi}(\tilde{F}_{V,Z})$ implies that $\psi(F_{V,Z}, \phi) = \psi(\tilde{F}_{V,Z}, \tilde{\phi})$.

In addition to identification, some of our results will analyze whether the implications of a model are refutable:

Definition 2. A model (\mathbb{F}, \mathbb{S}) is **falsifiable** if and only if $\bigcup_{\phi \in \mathbb{S}, F_{V,Z} \in \mathbb{F}} \phi(F_{V,Z}) \subsetneq \mathbb{H}$.

Just as identification of a model is necessary but not sufficient for the existence of a consistent estimator, falsifiability is necessary but not sufficient for the existence of a valid statistical

test (Berry and Haile, 2018). Both estimation and inference require additional statistical analysis. We leave such analyses for future research.

3 Examples

3.1 Single-Agent Problems

Example 1. Hedonic Demand Models. Consider the hedonic demand model in which consumer i 's indirect utility from purchasing good $k \in \{1, \dots, K\}$ is given by

$$\sum_j x_{kj} \beta_{ij} - p(x_k),$$

where $p(\cdot)$ is the pricing function, x_{kj} denotes the j -th characteristic of product k , and β_{ij} denotes the random co-efficients. This model fits our framework with $v_i = (\beta_{i1}, \dots, \beta_{iJ})$ and $t(\cdot) = p(\cdot)$.

This model is the hedonic demand model proposed in Gorman (1980) and Lancaster (1966) with the additional restriction that preferences are linear in characteristics and prices. Rosen (1974) proposed estimating such models by first estimating $p(\cdot)$ and then using data on purchases to recover the marginal willingness to pay for x_{kj} , β_{ij} , between x_{kj} . Bajari and Benkard (2005) incorporate price endogeneity into this framework by including an unobserved quality index $\xi_k \in \mathbb{R}$ into x_k and show how to identify the pricing function and the unobserved quality of each good in a first step. Our analysis takes this first step as given and applies to the identification of the marginal willingness to pay.

Example 2. Multinomial Choice with Exogenous Characteristics. Consider a discrete choice model (see McFadden, 1973; Train, 2009). An outcome $x \in \mathcal{X} = \{x \in \{0, 1\}^J : \sum_j x_j \leq 1\}$ denotes which option is chosen by a consumer. Let $t(x)$ denote the price of option x and let v_i denote the vector of utilities for the other attributes. The consumer's utility from picking any given $x \in \mathcal{X}$ is given by the form assumed in our model:

$$v_i \cdot x - t(x).$$

A large literature (e.g., Berry et al., 1995) focuses on solving the price endogeneity problem. This problem is particularly relevant for the consumer choice context when certain product attributes are unobserved. We do not address endogeneity of this form in our analysis,

assuming that the researcher is able to find a solution in a prior step or it is reasonable that endogeneity is not a concern. Thus, our results are relevant to choices in which unobserved product attributes can be controlled for by other means or if there is consumer-level price variation within the market (e.g., [Tebaldi et al., 2019](#)).

3.2 Mechanisms and Trading Games

Example 3. Single Unit, Independent Private Value (IPV) Auctions. Our model embeds standard IPV auctions that award the object to the highest bidder. In our notation, the action $a \in \mathbb{R}$ corresponds to a bid, v_i is agent i 's value for the object, $x_{\mathcal{A}}(a) \in [0, 1]$ denotes the probability of winning with a bid a , and $t_{\mathcal{A}}(a)$ denotes the expected payment. We can accommodate both first-price and all-pay auctions, amongst others. For example, in the first-price auction analyzed by [Guerre et al. \(2000\)](#), each agent chooses a bid a to maximize $(v - a)H(a)$, where $H(a)$ is the cumulative distribution function of the highest bid of the remaining bidders. In a Bayesian Nash Equilibrium, bidders have consistent beliefs about the bid distribution of opposing bidders and data from many independent and identical auctions identifies $H(a)$. The bid a parametrizes $x_{\mathcal{A}}(a) = H(a)$ and $t_{\mathcal{A}}(a) = aH(a)$.

We assume that all bids are observed. The identification of these auctions under weaker assumptions and fewer data requirements has been analyzed in [Athey and Haile \(2002\)](#).

Example 4. Mechanisms with Private Information, Independent Types, and Quasi-linear Utilities. Our model can incorporate mechanisms more general than single-unit IPV auctions. Consider a mechanism described by the allocation rule $\tilde{x}_i(a_i, a_{-i})$ and transfer function $\tilde{t}_i(a_i, a_{-i})$, where the set of actions \mathcal{A} coincides with the set of messages an agent can send. In incentive-compatible direct mechanisms, agents will truthfully reveal their types and the distribution of a_{-i} will coincide with the distribution of valuations. In indirect mechanisms, revealed preference arguments link valuations and messages. Moreover, the resulting allocation may involve ironing or pooling types. Pooling is particularly important if the space of messages is discrete, for example if there are minimum bid increments. An example with discrete messages includes scoring auctions in which bids may include binary services in addition to a continuous monetary amount (see [Aspelund and Russo, 2023](#), for example). Our approach will allow for both cases.

Example 5. Oligopoly Pricing. The identification of marginal costs in canonical models (e.g., [Berry et al., 1995](#)) is based on first identifying demand using cost shifters (see [Berry and Haile, 2014](#)), then assuming a Nash Equilibrium in simultaneous move price setting game. To

map this exercise to our notation, let a_i be the price chosen by firm i , v_i be the negative of the firm's marginal cost, $\tilde{x}(a_i, a_{-i})$ be the quantity sold by firm i , and $\tilde{t}(a_i, a_{-i})$ be the negative of the firm's revenue. Therefore, $x_{\mathcal{A}}(a_i; \mathcal{J}_i)$ and $t_{\mathcal{A}}(a_i; \mathcal{J}_i)$ are the expected quantities and the negative of expected revenues, respectively. Profit maximization implies that the firm maximizes

$$v_i \cdot x_{\mathcal{A}}(a_i; \mathcal{J}_i) - t_{\mathcal{A}}(a_i; \mathcal{J}_i).$$

We can allow for information sets that may or may not include the prices set by other firms by varying \mathcal{J}_i .

Instead of a price setting game, it is also straightforward to fit a quantity setting game into our model by interpreting a as quantities and setting $x_i(a_i; \mathcal{J}_i) = a_i$. The function $t_i(a_i; \mathcal{J}_i)$ still denotes the expected negative revenue.

Example 6. School Choice. Agarwal and Somaini (2018) consider a school assignment mechanism in which agents can submit rank order lists indexed by a .⁶ Assume that preferences are private information and each student knows the distribution from which the preferences of other students are drawn. In equilibrium, a student submitting the list a is assigned to one of J schools with probability vector $x_{\mathcal{A}}(a) \in \Delta^J \subseteq \mathbb{R}^J$, where $x_{\mathcal{A}}(a)$ is derived as in section 2.2. Let $d_i = (d_{i1}, \dots, d_{iJ})$ be the vector of distances of each school from student i , and let u_i denote the vector of indirect utilities from assignment into each school, net of distance. If preferences are linear in distance, then the expected utility from submitting list a is given by

$$(u_i - d_i) \cdot x_{\mathcal{A}}(a).$$

This model fits our framework by setting $t_{\mathcal{A}}(a; \mathcal{J}_i) = 0$ and $v_i = u_i - d_i$.

Example 7. Trading Games. Larsen and Zhang (2018) consider a trading game in which an agent can take a sequence of actions $a = (a_1, \dots, a_M)$. Following this sequence, the agent has a probability $x_{\mathcal{A}}(a) \in [0, 1]$ of engaging in a transaction and paying an expected (possibly negative) transfer $t_{\mathcal{A}}(a)$. The value of the trade for agent i is given by $v_i \in [\underline{v}_i, \bar{v}_i] \subseteq \mathbb{R}$. Therefore, the expected utility from the action a sequence is

$$v_i \cdot x_{\mathcal{A}}(a) - t_{\mathcal{A}}(a),$$

where the actions of other agents are integrated over as in section 2.2. This trading game is a one-dimensional special case of our model.

⁶We fix the priority type of the student and drop it from the notation for simplicity.

4 Revealed Preferences

4.1 Rationalizable Actions

Our first result derives a testable implication of the model and characterizes the consequences x that are optimal for some $v \in \mathbb{R}^J$.

Proposition 1. (i) *The outcome $x \in \mathcal{X}$ is optimal for type $v \in \mathbb{R}^J$ if and only if $t_{\mathcal{X}}(x) = t(x)$ and $v \in \partial t(x)$. (ii) *If $t_{\mathcal{X}}(\cdot)$ is convex and lower-semicontinuous, then for every $x \in \mathcal{X}$ there exists $v \in \mathbb{R}^J$ such that x is an optimal choice.**

Proof. Part (i). If: Let $v \in \partial t(x)$ and $t_{\mathcal{X}}(x) = t(x)$. Since $v \in \partial t(x)$, for all x' in the convex hull of \mathcal{X} , $v \cdot (x' - x) \leq t(x') - t(x)$. Observe that $t_{\mathcal{X}}(x) = t(x)$ and $t(x') \leq t_{\mathcal{X}}(x')$ for all $x' \in \mathcal{X}$. Therefore, we have that $v \cdot (x' - x) \leq t_{\mathcal{X}}(x') - t_{\mathcal{X}}(x)$ as required.

Only if: Suppose that $x \in \mathcal{X}$ is optimal for v . Towards a contradiction suppose that $t(x) < t_{\mathcal{X}}(x)$. Since $t(\cdot)$ is the convex hull of $t_{\mathcal{X}}(\cdot)$, there exist $x_j \in \mathcal{X}$ and weights $\alpha_j \geq 0$ such that $\sum \alpha_j = 1$, $\sum \alpha_j x_j = x$ and $\sum \alpha_j t_{\mathcal{X}}(x_j) = t(x)$. Therefore,

$$v \cdot x - t_{\mathcal{X}}(x) < \sum_j \alpha_j (v \cdot x_j - t_{\mathcal{X}}(x_j)).$$

Hence, it must be that there exists x_j such that $v \cdot x - t_{\mathcal{X}}(x) < v \cdot x_j - t_{\mathcal{X}}(x_j)$. This contradicts the assumption that x is optimal. Thus, $t_{\mathcal{X}}(x) \geq t(x)$, and by definition of $t(\cdot)$, $t(x) \leq t_{\mathcal{X}}(x)$. Therefore, $t_{\mathcal{X}}(x) = t(x)$.

To show that $v \in \partial t(x)$, assume towards another contradiction that $v \notin \partial t(x)$. That is, assume there exists x' in the convex hull of \mathcal{X} such that $v \cdot x - t(x) < v \cdot x' - t(x')$. Hence, there exist $x_j \in \mathcal{X}$ and weights $\alpha_j \geq 0$ such that $\sum \alpha_j = 1$ and $\sum \alpha_j (v \cdot x_j - t_{\mathcal{X}}(x_j)) = v \cdot x' - t(x')$. Since $t_{\mathcal{X}}(x) = t(x)$, it must be that there exists x_j such that $v \cdot x - t_{\mathcal{X}}(x) < v \cdot x_j - t_{\mathcal{X}}(x_j)$. This contradicts the assumption that x is optimal.

Part (ii). Under the maintained assumptions, $t_{\mathcal{X}}(x) = t(x)$ for all $x \in \mathcal{X}$ and $\partial t_{\mathcal{X}}(x) = \partial t(x)$. Moreover, $t(x)$ is continuous, implying that $\partial t(x)$ is non-empty. Part (i) implies the result. \square

Part (i) shows that linearity of payoffs in the model has testable implications. Specifically, outcomes x that result in payments larger than $t(x)$ are dominated. Figure 1 presents an illustration. Consider the point A and an agent with type v_A given by the slope of the tangent

through A . Lines parallel to this tangent are the indifference curves for an agent with type v_A , with utility increasing as t decreases. Therefore, the unique optimizer for this agent is x_A . In contrast, points in the neighborhood of x_B are not an optimal choice. An agent would rather pick either x_C or an outcome close to x_A . The outcome x_C is also optimal, but for more than one type. The slopes of the two dotted lines touching C , and convex combinations of them, are the types for which C is optimal.

That said, part (ii) shows that if the cost function $t_{\mathcal{X}}(\cdot)$ is convex and smooth, then essentially any choice can be rationalized. Convexity implies that $t_{\mathcal{X}}(x) = t(x)$ for all $x \in \mathcal{X}$, ruling out points like B in Figure 1. Lower semi-continuity rules out points like D and F in the graph of $t_{\mathcal{X}}(\cdot)$, illustrated in Figure 1. The outcome $x_D \notin \mathcal{X}$ and therefore it does not belong to the domain of $t_{\mathcal{X}}(\cdot)$ although it belongs to the domain of $t(\cdot)$. The outcome x_F is not optimal for any type because an agent can achieve an arbitrarily close outcome to x_F while discretely reducing the payment. Proposition 1 implies that x_F is not optimal because $\partial t(x)$ is not well-defined at this point. Together, convexity and lower semi-continuity imply that all solutions to the problem $\max_{x \in \mathcal{X}} v \cdot x - t_{\mathcal{X}}(x)$ also solve the problem $\max_{x \in \bar{\mathcal{X}}} v \cdot x - t(x)$.

The assumptions of convexity and semi-continuity are commonly imposed in the literature on identification. In first-price auctions (example 3), it is often assumed that $b + \frac{G(b)}{\partial G(b)/\partial g(b)}$, where b is a bid and $G(b)$ is the cumulative distribution function of the highest competitor bid, is increasing (see Assumption C2 in Guerre et al., 2000, for example). This assumption is identical to assuming that $t(x)$ is convex if it is twice differentiable. To see this, note that the first-derivative of $t(x) = xG^{-1}(x)$ is equal to

$$\nabla t(x) = G^{-1}(x) + \frac{x}{\partial G(G^{-1}(x))/\partial b}$$

and substitute $x = G(b)$. Similarly, in the hedonic demand model, if the price function, $t(x)$, were non-convex, then certain choices would be sub-optimal under the linear random co-efficients structure.

4.2 Identification with Rich Choice Environments

Our first identification result considers the case when $t(x)$ is differentiable. It is a corollary of Proposition 1, applied for each fixed value of $Z = z$.

Corollary 1. *Suppose Assumption 1 is satisfied and agent i chooses outcome $x_i \in \mathcal{X}$. If $t(\cdot)$ is differentiable at x_i , then v_i is identified. In particular, if $t(\cdot)$ is differentiable for all x in the support of X , then $F_{V,Z}$ is identified.*

The result shows that choosing $x \in \mathcal{X}$ uniquely determines the type of all agents making that choice. For this reason, we label such a choice environment as “rich.”

Two special cases of this result are identification of the hedonic choice models and first-price auctions. To see this, recall our calculation of the derivative of $t(x) = xG^{-1}(x)$ in first-price auctions (example 3). This derivative, calculated at $x = G(b)$, is precisely the virtual valuation of a bidder bidding b .

In the higher-dimensional case of hedonic demand models, the marginal cost of increasing x_j is $\partial t(x) / \partial x_j$. Therefore, if consumer i chooses a product or a consumption bundle described by x_i , then optimality implies that $\nabla t(x_i)$ is equal to the vector of marginal willingness to pay for each of the components, v_i .

4.3 Partial Identification with Coarse Choice Sets

The complementary case is when $t(x)$ is not differentiable. This case is particularly relevant when \mathcal{X} is not a connected subset of \mathbb{R}^J , for example, when the choice set is discrete. For any fixed value of the characteristic z , we observe the distribution of X that agents with that characteristic choose. Because the set $\partial t(x)$ is not a singleton, without any further restrictions, the identified set may not be a singleton.

Proposition 2. *Suppose Assumption 1 is satisfied and agent i chooses outcome $x_i \in \mathcal{X}$. The identified set of $F_{V,Z}$ is given by*

$$\mathcal{F}_I = \left\{ F_{V,Z} : \text{for all } z \text{ and } A \subseteq \mathcal{X}, \Pr(X \in A|z) \leq \int 1\{v \in \cup_{x \in A} \partial t(x)\} dF_{V|Z=z} \right\}.$$

Moreover, under the additional restriction that $F_{V|Z=z}$ admits a density $f_{V|Z=z}$, then (i) the identified set is given by

$$\mathcal{F}_I = \left\{ F_{V,Z} : \text{for all } z \text{ and } A \subseteq \mathcal{X}, \Pr(X \in A|z) = \int 1\{v \in \cup_{x \in A} \partial t(x)\} f_{V|Z=z}(v) dv \right\},$$

and (ii) \mathcal{F}_I is non-empty if and only if $P(X \in \mathcal{X}^\dagger|z) = 1$.

The result follows because $P(X = x|z) \leq \mathbb{P}(v \in \partial t(x)|z) = \int 1\{v \in \partial t(x)\} dF_{V|Z=z}$. The first inequality is a consequence of Proposition 1, and the second is definitional. Observe that the identified set is convex. This is the case because the restrictions on $F_{V|Z}$ are linear.

Under the restriction that $F_{V|Z=z}$ admits a density, $\Pr(V \in \mathcal{V}^\dagger|z) = 1$ because the complement of \mathcal{V}^\dagger relative to \mathbb{R}^J is a set of measure zero. Thus, ties occur with zero proba-

bility. Therefore, $\Pr(X \in \mathcal{X}^\dagger|z) = 1$. $\Pr(X \in A|z) \leq \int 1\{v \in \cup_{x \in A} \partial t(x)\} f_{V|Z=z}(v) dv = \int 1\{v \in \cup_{x \in A} \partial t(x) \cap \mathcal{V}^\dagger\} f_{V|Z=z}(v) dv \leq \Pr(X \in A \cap \mathcal{X}^\dagger|z) = \Pr(X \in A|z)$. The first inequality follows from Proposition 1, the first equality follows because $\Pr(V \in \mathcal{V}^\dagger|z) = 1$, the second inequality follows from the fact that $v \in \partial t(x) \cap \mathcal{V}^\dagger$ implies that x is the unique maximizer of profits for v . The last equality follows because $\Pr(X \in \mathcal{X}^\dagger|z) = 1$. That concludes the proof of (i). The “only if” part of (ii) is immediate from the arguments showing that $\Pr(X \in \mathcal{X}^\dagger|z) = 1$.

A one-dimensional version of this result for trading models is given in [Larsen and Zhang \(2018\)](#). Specifically, consider an agent, indexed by i , who picks actions that result in a probability of trade equal to x_i . Assume that the agent can also pick actions that instead yield either $x_i + \Delta_1$ or $x_i - \Delta_2$ for some $\Delta_1, \Delta_2 > 0$. Since x_i is optimal, it must be that

$$\frac{t(x_i) - t(x_i - \Delta_1)}{x_i - (x_i - \Delta_1)} \leq v_i \leq \frac{t(x_i) - t(x_i + \Delta_2)}{x_i - (x_i + \Delta_2)}.$$

Convexity of $t(\cdot)$ implies that the local deviations considered above are both necessary and sufficient for optimality.

Our result extends naturally to a higher-dimensional version of this problem. It also applies to other scenarios. As an example, consider a hedonic demand model with a discrete product set. In this context, our result identifies the set to which the vector representing consumers’ willingness to pay for specific product characteristics belongs.

In higher dimensions, when the support of actions, and consequently the support of x is discrete, the set $\partial t(x)$ at some of these points in the support has positive volume in R^J . However, if the action space includes both discrete and continuous elements, the set $\partial t(x)$ may neither be a singleton nor have positive volume in R^J . Instead, it forms a convex set with an affine hull $\text{aff}(\partial t(x))$, that has a dimension $\dim(\text{aff}(\partial t(x)))$ lying between 0 and J .

The argument suggests a two-step estimation strategy. The first step recovers the subgradients $\partial t(x)$, which may or may not be points. Computing the subgradients may first require estimating the function $t(x)$ in some contexts. The second step estimates the CDF $F_{V|Z}$. A common example is auctions, a case that we further develop in our application in Section 5.⁷

⁷[Aspelund and Russo \(2023\)](#) is a recent example that analyzes a scoring auction using this estimation procedure in a context that does not squarely fit into prior models.

4.4 Identification with Preference Shifters

We now show that even with coarse choice sets, variation in an observable shifter of preference, denoted z , can be used to point identify the model. We start with the case when z is a special regressor (Lewbel, 2000), that is, $v = u + z$, with $u \perp z$. Then, we generalize this result to the case when $v = u + g(z)$, where $g(\cdot)$ is a non-linear function.

The rest of this section makes the following assumptions on the distribution of u and the support of outcomes:

Assumption 2. (i) *The random variable u is independent of z and admits a density $f_U(u)$.*
(ii) *There exists a constant $k > 0$ such that $\exp(k|u|) f_U(u)$ is Lebesgue integrable.*

This assumption requires that the tails of u decline sufficiently rapidly. It is satisfied by most commonly used parametric forms, including multivariate normals and extreme-value distributions as well as finite mixtures of these distributions.

Assumption 3. *The convex hull of \mathcal{X} has a non-empty interior.*

This assumption allows to consider choice sets that are not “rich,” which is the focus of this subsection. In particular, if the set \mathcal{X} contains a finite number of options, the function $t(\cdot)$ will be differentiable everywhere except in the set $\mathcal{X}^\dagger \subseteq \mathcal{X}$. However, if the distribution of valuations admits a density, X belongs to \mathcal{X}^\dagger with probability one, which implies that all choices with positive probability result in a non-singleton $\partial t(x)$. Assumption 3 still requires some limited richness of the choice set. A non-empty interior implies that the dimension of \mathcal{X} and of $t(\cdot)$ is J . Thus, the choice of x is informative about every dimension of payoffs.

With these assumptions, we have the following result:

Theorem 1. *Suppose that $v = u + z$ and Assumptions 1, 2 and 3 are satisfied. If z has full support on \mathbb{R}^J , then $f_U(\cdot)$ is identified. Moreover, $f_U(\cdot)$ is identified if there exist a linear subspace $L \subseteq \mathbb{R}^J$ such that (i) for every x in the support of X , $t(x)$ is differentiable in every direction $d \in L$ and (ii) z has full support on L^\perp , the orthogonal complement of L .*

The condition of the theorem requires that z has sufficient variation in order to trace the tails of the distribution of u . This condition is similar to those used in other arguments that rely on special regressors.

Proof. Lemma 2 implies that there exists a set $A \in \bar{\mathcal{X}}$ that has strictly positive Lebesgue measure and is contained in a translated simplicial cone. Let $\mathcal{F}_k \subseteq \mathbb{L}^1(\mathbb{R}^J)$ be the space of functions that satisfy the integrability condition in Assumption 2 for a given value of k . Fix $k > 0$ such that $f_U \in \mathcal{F}_k$ for the true f_U . Define the operator $A : \mathcal{F}_k \rightarrow \mathbb{L}^\infty(\mathbb{R}^d)$ as

$$A[f](z) = \int 1\{u + z \in \cup_{x \in A} \partial t(x)\} f(u) du.$$

Note that $A[f_U](z) = \Pr(X \in A|z)$ for the true value of f_U . We will show that $A[f] = 0$ a.e. implies that $f = 0$ a.e. if $f \in \mathcal{F}_k$. This statement implies that f_U is identified because if $A[f_U](z) = \Pr(X \in A|z)$ for two candidate functions f_U , then linearity of the map A implies that the two functions have to be identical.

Towards a contradiction, suppose $A[f] = 0$ and f is nonzero on a set with positive Lebesgue measure. It is without loss to assume that $\cup_{x \in A} \partial t(x)$ is contained in a simplicial cone because if it is contained in $C + \{z_0\}$ where C is a simplicial cone, then we can re-define the operator above by replacing the argument of $A[f]$ with $z' = z - z_0$. Let $\mathcal{N}(C)$ be the normal cone to C . By the definition of \mathcal{F}_k , there exists $\lambda \in \text{int } \mathcal{N}(C)$ sufficiently small so that $\exp(2\pi u \cdot \lambda) f(u)$ is integrable for all $f \in \mathcal{F}_k$. Fix one such value of λ . Rewrite

$$\begin{aligned} A[f](z) &= \int 1\{u + z \in \cup_{x \in A} \partial t(x)\} f(u) du \\ &= \int 1\{u \in \cup_{x \in A} \partial t(x)\} f(u - z) du \\ &= \exp(2\pi z \cdot \lambda) \int 1\{u \in \cup_{x \in A} \partial t(x)\} \exp(-2\pi u \cdot \lambda) \exp(2\pi(u - z) \cdot \lambda) f(u - z) du. \end{aligned}$$

Since $\exp(2\pi z \cdot \lambda) > 0$ a.e., $A[f] = 0$ a.e. $\iff \hat{\chi}_{\partial t(x), \lambda}(\xi) \cdot \bar{\hat{f}}_\lambda(\xi) = 0$, where $\bar{\hat{f}}_\lambda$ is the conjugate of the Fourier transform of $f_\lambda(u) = \exp(2\pi u \cdot \lambda) f(u)$ and $\hat{\chi}_{\partial t(x), \lambda}$ is the Fourier transform of $\chi_{\partial t(x), \lambda} = 1\{u \in \partial t(x)\} \exp(-2\pi u \cdot \lambda)$. Since $\hat{f}_\lambda(\xi)$ is continuous, the set of values ξ where $\hat{f}_\lambda(\xi) \neq 0$ is open. Further, since $\|f_\lambda\|_1 > 0$, the support of $\hat{f}_\lambda(\xi)$ is non-empty. Therefore, there is a non-empty open set Ξ , such that $\hat{f}_\lambda(\xi) \neq 0$ for all $\xi \in \Xi$. Because $\hat{\chi}_{\partial t(x), \lambda}(\xi) \cdot \bar{\hat{f}}_\lambda(\xi) = 0$, it must be that for all $\xi \in \Xi$, $\hat{\chi}_{\partial t(x), \lambda}(\xi) = 0$. However, this conclusion contradicts Lemma 1 below, which shows that $\hat{\chi}_{\partial t(x), \lambda}$ is not zero on any open set $\Xi \subseteq \mathbb{R}^J$.

If $t(x)$ is differentiable in every direction $d \in L$ for every $x \in \bar{\mathcal{X}}$ then there is an orthonormal transformation T of outcomes and values so that the resulting $\tilde{t}(Tx) = t(x)$ is differentiable in the last d arguments, where d is the dimension of the subspace L . Thus, it is without loss of generality to assume that $t(x)$ is differentiable in the last d arguments. Therefore, the

subdifferential $\partial t(x)$ is the Cartesian product of a set $V_1 \subseteq \mathbb{R}^{J-d}$ and a point $v_2 \in \mathbb{R}^d$. Thus, $f_{U_1}(\cdot)$, the distribution of U_2 , which is defined as the last d elements of U , is identified from the observed distribution of the last d elements of $\partial t(x) - \{z\}$. Consider the distribution of U_1 , defined as the first $J - d$ elements of U , conditional on z and $U_2 = u_2$ in the support of U_2 . Let $z = [z_1, z_2]$, where z_1 has full support in \mathbb{R}^{J-d} and $z_2 \in \mathbb{R}^d$ that, for simplicity is assumed to take a constant value of zero.

Let $\bar{\mathcal{X}}_1 \subseteq \mathbb{R}^{J-d}$ be the projection of $\bar{\mathcal{X}} \subseteq \mathbb{R}^J$ onto its first $J - d$ dimensions. Consider the problem of an agent with type $v = [v_1, u_2]$ of choosing sequentially the optimal $x_1 \in \bar{\mathcal{X}}_1$ first followed by a choice of x_2 so that $[x_1, x_2] \in \bar{\mathcal{X}}$. Given a choice of x_1 , the agent's problem is to choose x_2 to maximize $u_2 x_2 - t_2(x_2|x_1)$, where $t_2(x_2|x_1) = t([x_1, x_2])$. The conjugate function $t_2^*(u_2|x_1)$ denotes the maximum utility that an agent of type v can achieve in the second step if it chooses x_1 in the first one. Now, consider the first problem. The agents solves $\max_{x_1 \in \bar{\mathcal{X}}_1} v_1 x_1 - t_1(x_1|u_2)$, where $t_1(x_1|u_2) = t_2^*(u_2|x_1)$. As a function of its first argument, $t_1(\cdot|u_2)$ is a lower semi-continuous convex function supported over a non-empty convex and compact set $\bar{\mathcal{X}}_1$ by Assumption 1. $f_{U_1|u_2}(\cdot)$ satisfies part (ii) of Assumption 2 for almost all u_2 , and \mathcal{X}_1 satisfies assumption 3. Thus, by the first part of this Theorem, $f_{U_1|u_2}(\cdot)$ is identified for almost all u_2 . Therefore, the joint distribution of U is identified. \square

The proof technique is based on Fourier deconvolution methods. Under Assumption 2, the distribution of v is obtained using a convolution of the distributions of u and z . The choice of a specific set A restricts the set of v to $\cup_{x \in A} \partial t(x)$. Therefore, by observing $P(X \in A|z)$ for various values of z , we obtain information about the distribution of u because $P(X \in A|z) = \int 1\{u + z \in \cup_{x \in A} \partial t(x)\} f(u) du$.

The result below states the key technical result in the argument, which is based on finding a vector λ such that $\lambda \cdot v > 0$ for all $v \in \partial t(x)$ and then working with the Fourier transform of the function $\chi_{\partial t(x), \lambda}(u) = 1\{u \in \cup_{x \in A} \partial t(x)\} \exp(-2\pi u \cdot \lambda)$:

Lemma 1. *Suppose $\partial t(x)$ has strictly positive Lebesgue measure and $\partial t(x)$ is contained in a (closed) simplicial cone. Then, there exists $\lambda \in \text{int } \mathcal{N}(A)$, with $|\lambda|$ arbitrarily small, such that*

$$\hat{\chi}_{C, \lambda}(\xi) = \int 1\{u \in C\} \exp(-2\pi u \cdot (i\xi + \lambda)) du$$

is not zero on any open set $\Xi \subseteq \mathbb{R}^J$.

The formal proof is presented in Appendix A. The argument has three parts. First, we show that $\hat{\chi}_{C, \lambda}(\xi)$, when viewed as a function with complex domain, is holomorphic in

a neighborhood of \mathbb{R}^J . This result is obtained by verifying the conditions necessary for differentiating under the integral sign when dealing with functions that have complex domain (Theorem 13.8.6(iii) in [Dieudonné, 1976](#)). Second, we apply Theorem 5 in [Shabat \(1992\)](#) which implies that if $\hat{\chi}_{C,\lambda}(\xi)$ is zero on an open subset of \mathbb{R}^J then it is zero everywhere. Third, we observe that this second conclusion contradicts the fact that $\chi_{C,\lambda}(u)$ is, up to scale, a density function.

The technical arguments are a significant generalization of Theorem A.2 in [Agarwal and Somaini \(2018\)](#). This previous result, which is special to a school choice model, dealt the case in which $\partial t(x)$ is a convex cone. When $\partial t(x)$ is a convex cone, the Fourier transform of $\chi_{\partial t(x),\lambda}(u)$ can be computed in closed form, circumventing the need for more general arguments.

A limitation of Theorem 1 is that it requires a linearly separable regressor z that is independent of u . Our next result applies relaxes this requirement by considering the model

$$v = u + g(z),$$

where $g : \mathbb{R}^J \rightarrow \mathbb{R}^J$ with the dimension of z , $d_z \geq J$. We use an argument inspired by [Allen and Rehbeck \(2017\)](#) to show that the function $g(\cdot)$ is identified under the following additional restrictions:

Assumption 4. (i) $J \geq 2$ and the function $g(z)$ is given by $(g_1(z_1), \dots, g_J(z_J))$. Moreover, each $g_j(\cdot)$ is differentiable at each point.

(ii) For any $l, k \in \{1, \dots, J\}$, the partial derivatives of each $\mathbb{E}[X_l|z]$ with respect to z_k exist, are continuous, and are non-zero for all z .

(iii) The support of Z is rectangular.

(iv) The expectation of U exists.

The main restrictions are in parts (i) and (ii). Part (i) assumes that each of the regressors z_j is component-specific and that there are at least two components. We will discuss the need for at least two components after presenting our main result. Extensions that allow for additional regressors, some of which are common, can be accommodated by following arguments in [Allen and Rehbeck \(2017\)](#). Part (ii) assumes that the expected value of the optimal x is smooth and the partial derivatives are non-zero. This assumption would be violated if a component of g were not globally either a strict complement or substitute with a

component of X . Since this quantity is observed in the data, this assumption is falsifiable.⁸

Parts (iii) and (iv) are technical regularity assumptions. Part (iii) allows us to move from knowing the derivatives of g to determining the function up to a location. Part (iv) is a weak condition that is implied, for example, by Assumption 2(ii).

Proposition 3. *Suppose that Assumptions 1 and 4 are satisfied and $u \perp z$. Then, g is identified on its support up to the location and scale normalizations $g(z_0) = 0$ and $\frac{\partial}{\partial z_j} g_j(z_0) \in \{-1, 1\}$ for some j , respectively.*

Proof. Define $v^*(g) = \mathbb{E}[\max_{x \in \mathcal{X}} x \cdot (u + g) - t(x) | g]$. This expectation exists because

$$\begin{aligned} \mathbb{E} \left[\max_{x \in \mathcal{X}} x \cdot (u + g) - t(x) \middle| g \right] &= \mathbb{E} [X^*(u + g) \cdot (u + g) - t(X^*(u + g)) | g] \\ &\leq \mathbb{E} [|X^*(u + g)| \cdot |u + g| + |t(X^*(u + g))| | g], \end{aligned}$$

which is finite since X^* belongs to compact-valued set, u is independent of g and has finite expectation, and $t(\cdot)$ is a continuous function. Since $X^*(v)$ is measurable, we have that

$$v^*(g) = \max_{X: \mathbb{R}^J \rightarrow \mathcal{X}} \int [X(u) \cdot (u + g) - t(X(u))] f_U(u) du.$$

The equality follows from setting $X(u) = X^*(u + g) \in \arg \max_{x \in \mathcal{X}} x \cdot (u + g) - t(x)$ to show a weak inequality in one direction, and the definition of $v^*(g)$ for the other. Re-writing, we get that

$$v^*(g) = \max_{X: \mathbb{R}^J \rightarrow \mathcal{X}} \left[g \cdot \int X(u) f_U(u) du + \int [X(u) \cdot u - t(X(u))] f_U(u) du \right].$$

Observe that the maximand is linear in g , and therefore equidifferentiable with respect to each g_j . Moreover, the partial derivative of the maximand with respect to each g_j is uniformly bounded because \mathcal{X} is compact. Therefore, by the generalized envelope theorem of [Milgrom and Segal \(2002\)](#) (see Theorem 3),

$$\nabla v^*(g) = \int X^*(u + g) f_U(u) du = \mathbb{E}[X^*(u + g(z)) | g(z) = g].$$

Differentiating, we get that

$$\frac{\partial \mathbb{E}[X_k | z]}{\partial z_l} = \frac{\partial \mathbb{E}[X_k^*(u + g(z)) | z]}{\partial z_l} = \partial_{l,k} v^*(g(z)) \frac{\partial g_l(z_l)}{\partial z_l},$$

⁸This assumption substitutes for the non-testable assumption in [Allen and Rehbeck \(2017\)](#) that the second-order cross partials of $\mathbb{E}[\max_{x \in \mathcal{X}} x \cdot (u + g(z)) - t(x) | z]$ are non-zero.

where the derivatives exist by Assumption 4(iii). Therefore, for any value of g and z , and any pair k and l , we can identify

$$\frac{\partial g_l(z_l)}{\partial z_l} / \frac{\partial g_k(z_k)}{\partial z_k}. \quad (3)$$

The rest of the proof uses the arguments in Corollary S.5.1 in Allen and Rehbeck (2017). First, we identify $\frac{\partial}{\partial z_j} g_j(z_0)$ up to scale. Optimality and independence of u and z implies that

$$\begin{aligned} (\mathbb{E}[X^*(u + g(z))|z] - \mathbb{E}[X^*(u + g(z'))|z']) \cdot g(z) &\geq (\mathbb{E}[X^*(u + g(z))|z] \cdot u - \mathbb{E}[X^*(u + g(z')) \cdot u|z']) \\ &\quad - (\mathbb{E}[t(X(u + g(z)))|z] - \mathbb{E}[t(X^*(u + g(z')))|z']). \end{aligned}$$

An identical expression holds in which the left hand side switches the roles of z and z' . Adding these two inequalities, we get that

$$(\mathbb{E}[X^*(u + g(z))|z] - \mathbb{E}[X^*(u + g(z'))|z']) \cdot (g(z) - g(z')) \geq 0.$$

If $\mathbb{E}[X^*(u + g(z))|z] \neq \mathbb{E}[X^*(u + g(z'))|z']$, then it must be that $g(z) \neq g(z')$ and

$$(\mathbb{E}[X^*(u + g(z))|z] - \mathbb{E}[X^*(u + g(z'))|z']) \cdot (g(z) - g(z')) > 0. \quad (4)$$

Now, consider a small change in the j -th component starting from z_0 . We know that

$$(\mathbb{E}[X^*(u + g(z_0))|z_0] - \mathbb{E}[X^*(u + g(z_0 + \Delta_j))|z_0 + \Delta_j]) \cdot (g(z_0) - g(z_0 + \Delta_j)) > 0,$$

where Δ_j is the j -th standard basis vector multiplied by a small $\Delta > 0$. Because regressors are dimension-specific, we have that the sign of $g_l(z_{0,j}) - g_l(z_{0,j} + \Delta)$ is identified for all $\Delta > 0$. Taking the limit of this difference divided by Δ as $\Delta \rightarrow 0$ implies that the sign of $\frac{\partial g_j(z_{0,j})}{\partial z_j}$ is identified for all j . The scale normalization implies that the partial derivatives of g are known at z_0 .

Identification of $\frac{\partial g_j(z_{0,j})}{\partial z_j}$ and the ratios in equation (3) implies that for all l , $\frac{\partial g_l(z)}{\partial z_l}$ is identified for all z in its support. To show that this is sufficient to identify $g(z)$ up to scale and location, assume that there are two functions $g(z)$ and $\tilde{g}(z)$ that have the same partial derivatives and let $\delta(z) = g(z) - \tilde{g}(z)$. Since $\nabla \delta(z) = \nabla g(z) - \nabla \tilde{g}(z) = 0$ and the support of z is rectangular, we have that δ is the constant function. The location normalization implies that g is identified. \square

The main observation is an envelope theorem argument to show that the gradient of $v^*(g)$ is equal to the expected value of X , which is observed. This argument is akin to Roy’s identity from consumer theory. The second derivatives then can be identified and can be used to learn about the derivatives of g .

There are two differences from [Allen and Rehbeck \(2017\)](#) that are worth noting. First, they study a more general model choice structure than ours but focus on the identification of $g(z)$ while treating the distribution of u as a nuisance parameter. We identify this distribution. Second, we shorten their proof substantially by side-stepping the “representative agent’s problem” they define.⁹

Theorems 1 and 3 immediately imply the following result:

Corollary 2. *Suppose the hypotheses of Theorem 1 and Proposition 3 are satisfied. If $v = u + g(z)$ and $g(z)$ has full support on \mathbb{R}^J , then $f_{V|Z}(v|z)$ is identified.*

5 Application: Combinatorial Auctions

5.1 Empirical Setting

In our empirical application, we analyze the combinatorial auction used to procure school lunches in Chile. The National Board for School Aid and Scholarships (JUNAEB) contracts with private catering companies to prepare meals and deliver them to schools in the 90 territorial units (TUs). The typical contract has a three-year duration and covers one or multiple TUs. [Epstein et al. \(2002\)](#) and [Kim et al. \(2014\)](#) provide additional institutional details; we continue to abbreviate the latter as KOW.

Each year, JUNAEB uses a combinatorial auction to procure meals for about a third of the TUs. A bid consists of a list of TUs, i.e., a package, and a per-meal quote. Bidders have to meet some technical and financial requirements to qualify for the auction. Depending on their qualifications, bidders may face restrictions on the packages they can bid, such as a maximum number of TUs or a maximum number of meals. The number of bids a single bidder can submit is limited to 10,000. JUNAEB determines the contract allocation by solving a linear program that minimizes the total cost and ensures contracts for all TUs.

⁹We found this direct proof to be simpler than applying the general conditions of Theorems 1 and 2 in [Allen and Rehbeck \(2017\)](#). A cost of our approach is that we forego the appealing interpretation of the observed quantity $\mathbb{E}[X|z]$ as the solution to the representative agent’s problem.

Table 1: Auction Characteristics

<i>Panel A: Auction environment</i>	
Total meals allocated (millions)	80.76
# TUs allocated	32
# bidders	20
# bids submitted	43,136
<i>Panel B: Awarded allocation</i>	
Payment (pesos per meal)	410.96
# winning bidders	9

Note: In these tables and elsewhere, the number of meals refers to the total number of meals served per year. Winning bidders are allocated three-year contracts.

We use the same dataset as in KOW. The primary dataset contains all the bids submitted to the auctions. Two supplementary datasets contain TU-specific geographical and demographic information and bidder-specific financial and technical ratings. The information in these two datasets allows us to construct all the relevant restrictions to the cost-minimization problem. We focus on the 2003 auction, allowing us to compare our results to KOW.

Table 1 describes the auction environment and the outcome. JUNAEB allocated 32 TUs, totaling about 81 million meals. There were twenty bidders, which together bid on over 43,000 packages. Nine of the bidders were included in the final allocation of 32 TUs. The average price bid was 423 pesos per meal. At the end of 2003, the exchange rate was 599 Chilean pesos per U.S. dollar (OECD, 2016). The average price bid in U.S. dollars was therefore 71 cents per meal.

Table 2 describes the bids in greater detail. Panel A shows that there is significant heterogeneity in the total number of meals across TUs, with the 25th percentile of TUs serving just over 2 million meals and the 75th percentile serving over 3 million meals. The number of package bids containing a TU also varies substantially, although each TU receives at least one bid from each bidder. Panel B shows that bidders face varying restrictions on the maximum number of meals they are allowed to bid and the maximum number of TUs they may be allocated. There is also significant heterogeneity in the number of packages that a bidder may bid on. However, the number of packages on which a bidder does place a bid is much lower than the number of feasible packages. Finally, panel C shows that the bid price per meal exhibits significant heterogeneity across firm-package pairs.

Table 2: TUs, Bidders, and Bids

	Min	25th per- centile	Median	75th per- centile	Max
<i>Panel A: TUs</i>					
Meals (millions)	1.64	2.07	2.40	3.09	3.79
# bidders on TU	20	20	20	20	20
# bids on packages containing TU	1,008	4,219	7,204	9,370	11,198
<i>Panel B: Bidders</i>					
Max meals allowed (millions)	25.92	35.11	37.77	38.77	50.02
Max TUs allowed	1	3	4	8	8
# TUs bid on	9	29	32	32	32
# packages bid on	17	505	1,164	3,679	9,157
# feasible packages	32	2,124	10,906	31,353	31,353
<i>Panel C: Bids</i>					
Price (pesos per meal)	299.59	401.92	421.41	440.35	690.65
# TUs in package	1	4	5	7	8

Note: The unit of observation is the TU in Panel A, the firm in Panel B, and the bid (i.e., the firm-package pair) in Panel C.

5.2 Empirical Model

5.2.1 Setup

Bidders and Packages: Our empirical setting is a special case of Example 4, Mechanisms with Private Information and Independent Types and Quasi-linear Utilities. Each TU, indexed by $k \in \{1, \dots, K\}$, must be allocated to one of the firms in the auction, indexed by $i \in \{1, \dots, I\}$. Let $j \in \{1, \dots, J\}$ index subsets of TUs, or **packages**, where $J = 2^K - 1$ is the number of possible nonempty packages. For each firm i , the outcome space \mathcal{X} is the space of vectors $x_i = \{x_{ij}\}_j$, where x_{ij} denotes firm i 's probability of being allocated package j . Denote firm i 's action with $a_i \in \mathcal{A}_i = \mathbb{R}_+^J$, a vector of bids indicating a price for each package. A firm may choose to bid only on a subset of packages. If a firm does not bid on a particular package, we adopt the convention that the bid on that package is infinite. Package bidding allows firms to express complementarities or substitutabilities in costs of supplying multiple geographical units.

Costs: Let c_{ij} denote firm i 's cost of supplying package j , which we collect into the vector $c_i = \{c_{ij}\}_j$. In the notation of section 2.1, bidder i 's valuation vector is $v_i = -c_i$. We assume independent private valuations conditional on observables z_i :

Assumption 5. *The cost vector of each bidder i , denoted c_i , is independent of the costs of the other bidders and is distributed according to $F_{C|z_i}$. Each bidder's valuation is private information. Bidder characteristics z_i and conditional cost distributions $F_{C|z_i}$ are common knowledge.*

Types: We want to estimate the conditional valuation distribution $F_{C|z_i}$, but the high-dimensional nature of this problem makes estimation challenging. The spaces of outcomes, actions, and valuations all have dimension equal to the number of possible packages, $J = 2^K - 1$, which grows exponentially in the number of TUs, K . This curse of dimensionality makes calculating $\partial t(\cdot)$ challenging.

We reduce the dimensionality of the problem by assuming that firms' cost vectors $c_i \in \mathbb{R}^J$ (and therefore their valuation vectors) are a linear function of lower-dimensional types $\gamma_i \in \mathbb{R}^L$:

Assumption 6. *Costs satisfy $c_i = M\gamma_i$ for all firms i , where M is a known $J \times L$ matrix with $L \ll J$.*

Our objective is therefore to estimate the distribution of types $F_{\Gamma|z_i}$ given each bidder i 's observables z_i .

This approach contrasts with that in KOW, which solves the dimensionality problem by assuming that the markups that a firm bids, which are endogenous, are a low-dimensional function of package characteristics. Instead, we directly place restrictions on the distribution of firm costs as opposed to markups. These restrictions will allow us to ease the computation of types consistent with optimal bidding. As we will see below, these restrictions will reduce dimensionality when computing $\partial t(\cdot)$.

5.2.2 Mechanism and Equilibrium

Mechanism: In the notation of section 2.2, the Chilean combinatorial auction can be represented by an allocation rule $\tilde{x}_i(a_i, a_{-i})$ and a transfer function $\tilde{t}_i(a_i, a_{-i})$ for each firm i . Winning firms are paid the amount of their winning bids: $\tilde{t}_i(a_i, a_{-i}) = -a_i \cdot \tilde{x}_i(a_i, a_{-i})$.

The winning allocation is determined by finding the lowest total price at which all TUs can be served. This problem, known as the winner determination problem, can be written as an integer program. Stack the allocation rules for each firm into the function $\tilde{x} : \prod_i \mathcal{A}_i \rightarrow \prod_i \mathcal{X}_i$. The winner determination problem solves

$$\begin{aligned} \tilde{x}(a_1, \dots, a_I) \in \operatorname{argmin}_x \sum_i \sum_j x_{ij} a_{ij} \\ \text{s.t. } x \in \tilde{\mathcal{X}}, \end{aligned} \quad (5)$$

where $x = \{x_{ij}\}_{i,j}$ and $\tilde{\mathcal{X}} \subseteq \mathcal{X}^I$ is a set of allowable allocations. The constraints on $\tilde{\mathcal{X}}$ imposed by JUNAEB are (i) each TU k must be served – for each k , $\sum_i \sum_{j:k \in j} x_{ij} \geq 1$; (ii) no firm wins more than one package – for each i , $\sum_j x_{ij} \leq 1$; and (iii) a set of constraints on each firm indicating limits on their market share.¹⁰ These constraints, including the third set, are linear in x_{ij} . Generically, ties in the winner determination problem happen with probability zero.

Equilibrium: Before submitting her bid, a bidder observes her own type and characteristics, as well as her competitors' characteristics. A (mixed) strategy is a function $\sigma_i(\cdot; z_i, z_{-i}) : \mathbb{R}^L \rightarrow \Delta \mathcal{A}_i$ where the domain is the bidder's type γ_i and the range is a distribution over bid vectors. Let $F_{A_{-i}|z_i, z_{-i}}(a_{-i})$ denote the distribution of firm i 's opponents' bids conditional on

¹⁰The full set of feasibility constraints is presented in appendix B.1.

observables. A change of variables in equation (1) yields firm i 's expected outcome function

$$x_{\mathcal{A}}(a_i) = \int \tilde{x}_i(a_i, a_{-i}) dF_{A_{-i}|z_i, z_{-i}}(a_{-i}) \quad (6)$$

and (pay-as-bid) expected payment function

$$t_{\mathcal{A}}(a_i) = -a_i \cdot x_{\mathcal{A}}(a_i). \quad (7)$$

Assumption 7. *Bidder strategies follow a Bayesian Nash equilibrium.*

The strategy profile $(\sigma_1^*, \dots, \sigma_I^*)$ is a Bayesian Nash equilibrium if for each firm i and bid vector a_i with $\sigma_{i, a_i}^* > 0$,

$$\begin{aligned} a_i &\in \operatorname{argmax}_{a \in \mathcal{A}} -c_i \cdot x_{\mathcal{A}}(a_i) - t_{\mathcal{A}}(a_i) \\ &= \operatorname{argmax}_{a \in \mathcal{A}} (-c_i + a) \cdot x_{\mathcal{A}}(a) \\ &= \operatorname{argmax}_{a \in \mathcal{A}} (a - M\gamma_i) \cdot x_{\mathcal{A}}(a) \end{aligned} \quad (8)$$

As described in section 3.2, this setup bears close resemblance to GPV in the case of a single-unit auction.

5.2.3 Revealed Preference Type Bounds

The optimality condition (8) forms the basis for our estimation strategy. Given that bidder i submits a_i , define the **type bounds** $\mathcal{G}_i \subseteq \mathbb{R}^L$, the set of types γ_i for which $a_i \in \operatorname{argmax}_{a \in \mathcal{A}} (a - M\gamma_i) \cdot x_{\mathcal{A}}(a)$. Then, under Assumption 6, $-c_i$ is in the subdifferential $\partial t(x)$ if and only if $\gamma_i \in \mathcal{G}_i$. That is, \mathcal{G}_i is the lower-dimensional analogue in type space to the subdifferential.

Equivalently, \mathcal{G}_i is the set of types such that for all deviating bid vectors a'_i ,

$$M\gamma_i \cdot [x_{\mathcal{A}}(a_i) - x_{\mathcal{A}}(a'_i)] \leq a_i \cdot x_{\mathcal{A}}(a_i) - a'_i \cdot x_{\mathcal{A}}(a'_i). \quad (9)$$

For each deviation a'_i , the revealed preference inequality (9) specifies a halfspace in the domain of types. The type bounds \mathcal{G}_i are the intersection of these halfspaces for all possible deviations, forming a convex polyhedral set.

For each firm i , the empirical objects we target are: (i) the distribution $F_{A_{-i}|z_i, z_{-i}}$ of opponents' bids, (ii) the type bounds \mathcal{G}_i , and (iii) the conditional valuation distribution $F_{\Gamma_i|z_i}$.

The estimated bid distribution is used to evaluate the allocation function $x_{\mathcal{A}}(\cdot)$ via simulation of equation (6). The type bounds are obtained from the revealed preference inequality (9). These bounds are inputs to estimation of the conditional type distribution, our ultimate object of interest. We now describe our parameterizations of the bid and type distributions; the latter parameterization is only required for a subset of our empirical results.

5.2.4 Bid Distribution

Firm bids depend on two package characteristics. A package’s **volume** is its annual number of meals to be supplied, in millions. Abusing notation, we also use q_k to denote the volume of constituent TU k , so that $q_j = \sum_{k \in j} q_k$. Our **density** measure captures the geographic concentration of TUs within a package and takes values in the unit interval.¹¹ We discretize these variables into volume and density bins $\{\mathcal{Q}_0, \mathcal{Q}_1, \dots, \mathcal{Q}_{L^{\text{volume}}}\}$ and $\{\mathcal{D}_0, \mathcal{D}_1, \dots, \mathcal{D}_{L^{\text{density}}}\}$, respectively.

Assume that the distribution of $F_{A_i|z_i, z_{-i}}$ of firm i ’s per-meal equilibrium bids (from the perspective of i ’s competitors) is given by

$$\frac{a_{ij}}{q_j} = \sum_{k \in j} \frac{q_k}{q_j} \tilde{\beta}_{ik}^{\text{TU}} + \sum_{l=1}^{L^{\text{volume}}} \mathbf{1}\{q_j \in \mathcal{Q}_l\} \beta_{l, \text{size}_i}^{\text{volume}} + \sum_{l=1}^{L^{\text{density}}} \mathbf{1}\{d_j \in \mathcal{D}_l\} \beta_{l, \text{size}_i}^{\text{density}} + \epsilon_{ij}, \quad (10)$$

where size_i is a firm-specific categorical variable taking values “small” and “large.” The first right-hand side term is a TU volume-weighted sum of **base prices** $\tilde{\beta}_{ik}^{\text{TU}}$ charged for each constituent TU k . The base prices are distributed as

$$\tilde{\beta}_{ik}^{\text{TU}} = \beta_k^{\text{TU}} + \beta^{\text{incumb}} \text{incumb}_{ik} + \omega_{ik}, \quad (11)$$

where incumb_{ik} is an indicator for whether firm i is the incumbent supplier in TU k .¹² The base prices depend on (across-firm) means $(\beta_k^{\text{TU}})_k$ and an incumbency shifter β^{incumb} ,

¹¹We define the density of package j as $d_j = \sum_r \left(\frac{\sum_{k \in j \cap r} q_k}{q_j} \right)^2$, where the TUs have been partitioned into regions indexed by $r \in \{1, \dots, R\}$. This density measure can be interpreted as the probability that two randomly selected meals from package j will come from the same region. Packages with higher densities have TUs which are more geographically concentrated. Notice that by definition, single-TU packages are contained in a single region, and thus have densities of one. The opposite extreme is a geographically diffuse package with many TUs, each in its own region and with roughly the same number of meals; this package has a low density.

¹²The third set of allocation constraints in the winner determination problem includes constraints on how many TUs each firm is allowed to win. No firm is allowed to win more than 8 TUs. We call a firm small if it is allowed to win at most 6 TUs and large if it is allowed to win 7 or 8 TUs.

as well as a normally distributed shock $\omega_i = (\omega_{i1}, \dots, \omega_{iK}) \sim N(0, \Sigma_\omega)$. The next two terms are **bid adjustments** due to volume and density, where the smallest volume bin and largest density bin are omitted. The bid adjustment amounts $\beta^{\text{volume}} = \{\beta_{l,\text{size}}^{\text{volume}}\}_{l,\text{size}}$ and $\beta^{\text{density}} = \{\beta_{l,\text{size}}^{\text{density}}\}_{l,\text{size}}$ are different for small and large firms. The final term is an idiosyncratic error ϵ_{ij} ; it is normally distributed with mean zero and variance $\sigma_{\epsilon,|j|}^2$ depending on the number of TUs in the package.¹³

5.2.5 Type Distribution

Combinatorial auctions attempt to take advantage of cost complementarities across units. By allowing package bidding, they seek to mitigate the exposure problem present in simultaneous or sequential single-unit auctions.¹⁴ The effectiveness of package bidding depends on the magnitude of and across-firm heterogeneity in these complementarities, as well as on the degree of pass-through from cost complementarities to bid discounts.

We allow for complementarities in costs for supplying package j to depend on the volume q_j and the density d_j of the package. The map M from types to costs is specified as follows: firm i 's per-meal cost of supplying package j is

$$\frac{c_{ij}}{q_j} = \sum_{k \in j} \frac{q_k}{q_j} \gamma_{ik}^{\text{TU}} + \sum_{l=1}^{L^{\text{volume}}} \mathbf{1}\{q_j \in \mathcal{Q}_l\} \gamma_{il}^{\text{volume}} + \sum_{l=1}^{L^{\text{density}}} \mathbf{1}\{d_j \in \mathcal{D}_l\} \gamma_{il}^{\text{density}} \quad (12)$$

This cost is the sum of (i) the TU volume-weighted average of the costs γ_{ik}^{TU} of supplying each TU k in the package, (ii) cost complementarities or substitutabilities due to economies or diseconomies of scale, and (iii) the same due to economies or diseconomies of density. Equation (12) above parallels equation (10), which parameterizes the distribution of firm i 's bid on package j .

Firm i 's type is therefore $\gamma_i = \left((\gamma_{il}^{\text{volume}})_l, (\gamma_{il}^{\text{density}})_l, (\gamma_{ik}^{\text{TU}})_k \right) \in \mathbb{R}^L$.

For a subset of results, we will parametrize the conditional type distribution $F_{\Gamma_i|z_i}$ as a function of firm size and incumbency. Conditional on these observables, the type distribution is multivariate normal with variance-covariance matrix Σ_ν . The mean values of the volume

¹³Our bid distribution parameterization largely parallels that of KOW. We define package density differently and allow the variance-covariance matrix Σ_ω to be unrestricted.

¹⁴As an extreme example, suppose a firm views TUs A and B as perfect complements: its cost of supplying the two-unit package is 400 pesos per meal, while its cost of supplying either of the single-unit packages is infinite. Without package bidding, the firm always faces the risk of being allocated only one of the TUs, a prospect that it finds infinitely undesirable. With package bidding, the firm can choose to bid on the two-unit package but not the single-unit packages, so that it will never be allocated only a single TU.

and density components are $\mu_{\text{size}_i}^{\text{volume}}$ and $\mu_{\text{size}_i}^{\text{density}}$ respectively, again potentially differing by firm size. Mean TU costs are given by $\mu^{\text{TU}} + \mu^{\text{incumb}} \text{incumb}_i$, the sum of a common mean vector and an incumbency shifter.

5.3 Estimation Procedure and Results

The estimation procedure mimics the identification approach. First, we construct the outcome function $x_{\mathcal{A}}(a_i)$ and the expected payment function $t_{\mathcal{A}}(a_i)$ by estimating the bid distribution $F_{A_i|z_i, z_{-i}}$. Next, we estimate the type bounds \mathcal{G}_i using the revealed preference inequality (9). Finally, we estimate the conditional type distribution $F_{\Gamma_i|z_i}$.

5.3.1 Bid Distribution


We estimate the bid distribution parameters $\theta_a \equiv (\theta_{a,1}, \theta_{a,2})$ in two stages. First, we estimate $\theta_{a,1} = (\beta^{\text{volume}}, \beta^{\text{density}}, (\sigma_{\epsilon,n}^2)_n)$ and TU base prices $(\tilde{\beta}_{ik})_{ik}$ from equation (10) using feasible generalized least squares (FGLS). Second, we estimate $\theta_{a,2} = ((\beta_k^{\text{TU}})_k, \beta^{\text{incumb}}, \Sigma_{\omega})$ from equation (11) again using FGLS, taking the first-stage $(\tilde{\beta}_{ik})_{ik}$ estimates as data.

Across all firms and all TUs, the mean TU base price is about 464 pesos per meal, though this masks considerable variation, particularly across TUs. Figure 2 plots the distribution of TU base prices. The mean base price for TU k , denoted β_k^{TU} in equation (11), ranges from 356 to 607 pesos per meal. Within a TU, firms' base prices are correlated. However, we estimate that the incumbent firm in a TU submits bids for that TU that are 17.62 pesos per meal lower on average than those of its non-incumbent competitors, suggesting a potential incumbency cost advantage.

We find evidence of significant downward bid adjustments due to volume and density. Bid adjustments become more negative as package volume and density increase. We estimate that the prices bid on the largest (highest-volume) packages we observe are, on average, 25.7 to 29.3 pesos lower than those on the smallest packages. Likewise, we estimate that average bids on low-density packages are 1.1 to 5.6 pesos higher than those on high-density ones. Figure 3 shows the estimated bid adjustment functions for small and large firms. Small firms offer smaller discounts for package volume than large firms do. However, the pattern is reversed for density: it is small firms that offer the greater discounts for package density.


The estimated bid distribution fits the data well; we compare it to the observed bid distribution in figure B.1.

Figure 2: Distribution of Firms' TU Base Prices



Note: The blue boxplots show the distribution of non-incumbent firms' base prices charged for each TU, denoted $\tilde{\beta}_{ik}^{TU}$ in equation (10). Boxes show the 25th percentile, median, and 75th percentile. Lower whiskers extend to the lowest observed data point that is within a distance of 1.5 times the interquartile range (IQR) from the 25th percentile. Likewise, upper whiskers extend to the highest observed data point within 1.5 times the IQR from the 75th percentile. Blue dots indicate outliers. Each orange dot indicates the base price charged by the incumbent firm serving that TU.

Figure 3: Estimated Bid Adjustment Functions



Note: Figures show estimated bid adjustments due to economies of volume and density. Shaded areas show 95 percent confidence intervals computed using the standard errors of the estimated coefficients. The lowest volume bin and the highest density bin are omitted base categories, so bid adjustments are in comparison to low-volume, high-density packages.

5.3.2 Outcome and Payment Functions

Given bid distribution parameter estimates $\hat{\theta}_a$, we estimate the outcome function $x_{\mathcal{A}}(\cdot)$ and payment function $t_{\mathcal{A}}(\cdot)$ via simulation of equations (6) and (7), respectively. For each firm i , we use $S_a = 1,000$ competitor bids a_{-i} from the distribution $F_{A_{-i}|z_i, z_{-i}}(\cdot; \hat{\theta}_a)$.


Firms vary substantially in their probabilities of being included in the final allocation. Two firms are estimated to win with zero probability at their submitted bids, while the win probabilities of another four firms are estimated to be greater than 95 percent. As expected, firms' expected revenues $-t_{\mathcal{A}}(a_i)$ increase in their win probabilities, as seen in figure 4. Firms who submit lower average bids tend to have a greater probability of winning a package. Although not shown, win probabilities are, perhaps surprisingly, not very correlated with the *number* of bids submitted.

5.3.3 Type Bounds

Next, for each firm i , we estimate the set of types \mathcal{G}_i that satisfy the revealed preference inequality (9). There are two challenges in this exercise.

First, there are a large number of potential deviations and it is computationally costly to check all of them. Simulating the expected allocation function $x_{\mathcal{A}}(\cdot)$ requires repeated solutions

Figure 4: Bidders' Average Prices, Win Probabilities, and Expected Revenues



Note: Each circle represents a bidder. The marker size is proportional to the number of bids submitted by that firm. The horizontal axis indicates the firm's probability of winning any package—that is, the sum of the elements of $x_a(a_i)$, the expected allocation vector. In the left panel, the vertical axis indicates the firm's average per-meal price bid. In the right panel, the vertical axis indicates the firm's expected revenue, $t_a(a_i)$. These figures omit two firms which are estimated to have zero win probability.

of the winner determination problem at each of the deviating bid vectors and each of the $S_a = 1,000$ draws from the estimated bid distribution. Checking many deviations also results in a complex description of the set \mathcal{G}_i .¹⁵

The second challenge is that the expected allocation function $x_{\mathcal{A}}(\cdot)$ is estimated with simulation error. Considering larger deviations – to bid vectors farther away from the observed a_i – allows the differences $x_{\mathcal{A}}(a_i) - x_{\mathcal{A}}(a'_i)$ and $a_i \cdot x_{\mathcal{A}}(a_i) - a'_i \cdot x_{\mathcal{A}}(a'_i)$ in inequality (9) to be estimated more precisely, decreasing the variance but at the cost of a larger than necessary set \mathcal{G}_i .

To address these challenges, we estimate \mathcal{G}_i by checking the inequality in equation (9) for a subset of deviations $a'_i \in \tilde{A}_i \subset \mathcal{A}_i$. Observe that this results in a larger set of types that contains the true set \mathcal{G}_i . We consider two types of deviating bid vectors a'_i . In each deviation, we change the firm's bid on a single package j and hold constant its bids on other packages, $a'_{i,-j} = a_{i,-j}$. The two types of deviations are:

1. **Downward deviations:** For each package j in a subset $\tilde{\mathcal{J}}_i$ of possible packages, decrease the bid on package j by 50 pesos per meal.

¹⁵As the number of halfspaces grows, polyhedral operations like checking set membership (i.e., checking whether a given type vector lies in \mathcal{G}_i), projection onto subspaces, and elimination of dimensions become more difficult.

2. **Grid deviations:** For each single-unit package j (including j that we do not observe the firm bidding on in the data), deviate to bidding 300, 350, 400, 450, and 500 pesos per meal on j . This grid of per-meal bids roughly spans the range of observed prices bid.

We impose two additional constraints on the sets \mathcal{G}_i : (i) costs are below bids ($M\gamma_i \leq a_i$) and (ii) TU costs are nonnegative ($\gamma_i^{\text{TU}} \geq 0$). These inequalities don't require simulating the allocation function $x_{\mathcal{A}}(\cdot)$ so they are easier to compute.

The revealed preference inequalities and the two additional constraints yield a convex polyhedral set, which is in most cases bounded. Each downward deviation implies a lower bound on some linear combination of type dimensions: if firm i did not find it optimal to decrease its bid on package j by 50 pesos per meal, then its cost of supplying j must be *at least* something. Constraint (i) generates upper bounds on linear combinations of type dimensions.¹⁶ The grid deviations and constraint (ii) ensure that TU costs are bounded even if there are TUs that firm i does not bid on in the data. The downward deviation packages $\tilde{\mathcal{J}}_i$ are chosen to generate independent variation in each of the type dimensions: volume economies, density economies, and TU costs. We describe the construction of this set in appendix B.3.


Figure 5 shows type bound estimates for an example firm. There are four volume economies (the lowest volume bin is omitted), two density economies (the highest density bin is omitted), and 32 TU costs. Each black whisker in figure 5 plots a projection of the polyhedral set \mathcal{G}_i onto a single dimension. This example firm is not allowed to win any packages involving volume bins 3 and 4 or density bin 2, so we are unable to bound its realizations of those dimensions. The estimated bounds suggest that this firm has economies of volume, though we cannot rule out very small diseconomies of volume.¹⁷ Likewise, this firm exhibits either small economies or small diseconomies of density. Its TU costs are often but not always bounded from above by both its prices bid on single-unit packages and the TU base prices it bids.¹⁸

¹⁶In principle, we could also obtain upper bounds from the revealed preference inequalities implied by upward deviations (increasing per-meal bids on packages in $\tilde{\mathcal{J}}_i$). However, computing these requires simulating $x_{\mathcal{A}}(\cdot)$ and the resulting upper bounds are not that informative conditional on the bid upper bounds. As a result, we do not include upward deviations in $\tilde{\mathcal{A}}_i$.

¹⁷Higher values for the type dimensions $\gamma_{il}^{\text{volume}}$ and $\gamma_{il}^{\text{density}}$ increase the per-meal package cost in equation (12). Hence, a more negative value for $\gamma_{il}^{\text{volume}}$ indicates a larger economy of volume, while a more positive value indicates a larger *diseconomy* of volume.

¹⁸This does not contradict package costs being weakly below package bids because the cost of a single-unit package may also be a function of volume and/or density economies. For example, TU 401 has enough meals to fall in the first (non-omitted) volume bin. If j is the single-unit package containing only TU 401, then firm i 's per-meal cost of supplying package j is $\frac{c_{ij}}{q_j} = \gamma_{i1}^{\text{volume}} + \gamma_{i,401}^{\text{TU}} \leq \frac{a_{ij}}{q_{ij}}$. Since for this firm, we can't rule out negative values of $\gamma_{i1}^{\text{volume}}$, we also can't rule out some TU costs $\gamma_{i,401}^{\text{TU}}$ above the per-meal price bid $\frac{a_{ij}}{q_j}$.

Figure 5: Example Firm Type Estimates



Note: This figure plots type estimates for an example firm i . For each of the TU costs γ_{ik}^{TU} , the blue dot indicates firm i 's bid on the single-unit package containing only TU k . The orange dot indicates firm i 's base price bid on TU k , denoted $\tilde{\beta}_{ik}$. Each black whisker is the projection of the type bounds \mathcal{G}_i onto a single dimension. The left green violin plots show the marginals of the proposal distribution G_i . The right purple violin plots show the marginals of the estimated type distribution $F_{\Gamma_i|z_i}(\cdot | \hat{\theta}_\gamma)$. Both the proposal distribution and the estimated distribution are conditional on the type bounds \mathcal{G}_i .

5.3.4 Type Distribution

Finally, we estimate the parameters $\theta_\gamma = (\mu^{\text{volume}}, \mu^{\text{density}}, (\mu_k^{\text{TU}})_k, \mu^{\text{incumb}}, \Sigma_\nu)$ of the conditional type distribution $F_{\Gamma_i|z_i}$. The likelihood of θ_γ given firm i 's type bounds \mathcal{G}_i is

$$L(\theta_\gamma | \mathcal{G}_i) = \int 1\{\gamma_i \in \mathcal{G}_i\} dF_{\Gamma_i|z_i}(\gamma_i; \theta_\gamma). \quad (13)$$

We will simulate this likelihood because it need not yield a closed form solution. In principle, it suffices to count how many draws from $F_{\Gamma_i|z_i}(\cdot | \theta_\gamma)$ lie in the set \mathcal{G}_i , though a precise estimate of a small likelihood – if $F_{\Gamma_i|z_i}(\cdot | \theta_\gamma)$ puts little mass on \mathcal{G}_i – may require a large number of draws. We instead evaluate the likelihood at each candidate θ_γ using importance sampling (Ackerberg, 2009), using draws from a proposal distribution G_i which we reweight by their likelihood ratios:

$$\begin{aligned} L(\theta_\gamma | \mathcal{G}_i) &= \int 1\{x \in \mathcal{G}_i\} \frac{f_{\Gamma_i|z_i}(x; \theta_\gamma)}{g_i(x)} dG_i(x) \\ &\approx \frac{1}{S} \sum_{s=1}^S 1\{x_s \in \mathcal{G}_i\} \frac{f_{\Gamma_i|z_i}(x_s; \theta_\gamma)}{g_i(x_s)}, \end{aligned} \quad (14)$$

where $\{x_s\}_1^S$ denotes S simulated draws from the proposal distribution G_i . In practice, drawing from the unconditional proposal distribution is computationally burdensome because the high-dimensional nature of this problem implies that obtaining a draw $x_s \in \mathcal{G}_i$ has low probability. To solve this issue, we take advantage of the fact that the bounds \mathcal{G}_i have been previously computed and draw x_s from a set that is either equal to or slightly larger than \mathcal{G}_i .

Finally, we estimate θ_γ using a Metropolis-Hastings algorithm with the simulated version of $L(\theta_\gamma | \mathcal{G}_i)$.¹⁹ We describe our importance sampling and Metropolis-Hastings procedures in greater detail in appendix B.4.

The green violin plots in figure 5 show, for the same example firm i as above, the marginals of the proposal distribution G_i .²⁰ The purple violin plots show the marginals of the estimated type distribution $F_{\Gamma_i|z_i}(\cdot | \hat{\theta}_\gamma)$. Both distributions are shown conditional on the example

¹⁹This algorithm generates a Markov chain which converges to the posterior distribution. We take the posterior mean parameters as our point estimate $\hat{\theta}_\gamma$. We use the posterior mean rather than the mode because the former is more robustly estimated when the parameter space is high-dimensional and the posterior distribution has potentially many local maxima. We use a flat prior, so the posterior probability of each candidate θ_γ equals the likelihood. The posterior mode is therefore equivalent to the maximum likelihood estimator.

²⁰Tables B.4 and B.5 report summary statistics for the sampled posterior draws for all firms, and Figure B.2 compares the distributions of firms' TU base prices β_{ik} and TU costs γ_{ik}^{TU} .

firm's bids – that is, conditional on \mathcal{G}_i .²¹

For this example firm, our estimated type bounds do not rule out economies of volume of up to 118 pesos per meal, but nor do they indicate that such large volume economies (i.e., such negative values for the volume economy dimensions) are likely. Recall that the type bound estimates (black whiskers) plotted in figure 5 are one-dimensional projections of the polyhedral set \mathcal{G}_i . Thus, figure 5 shows that there exists a vector of the other type dimensions at which we can rationalize volume economies of 118 pesos per meal. However, the proposal distribution draws place little mass in this region. Since we use a conditional proposal distribution that is uniform on \mathcal{G}_i , this indicates that \mathcal{G}_i also has little volume in that region. The estimated type distribution places even less mass in that area; it also has little mass on values of the TU costs above the firm's per-meal prices bid.²²

5.4 Markups and Efficiency

5.4.1 CA Markups

Our first set of markup estimates uses only the estimates of the type bounds \mathcal{G}_i and does not require the additional parameterization of the type distribution. For each bid a_{ij} submitted by firm i on package j , we compute the minimum and maximum markups $\frac{a_{ij}-c_{ij}}{a_{ij}} = \frac{a_{ij}-M_j\gamma_i}{a_{ij}}$ that are consistent with the type bounds \mathcal{G}_i . The distributions of these lower and upper bounds across bids are shown in blue and orange, respectively, in Figure 6. We bound the aggregate markup on awarded bids from below by 0.5 percent of the total payment to winning bidders and from above by 31.6 percent.


Second, we compute expected markups using the estimated type distribution $F_{\Gamma_i|z_i}(\cdot | \hat{\theta}_\gamma)$ conditional on the type bounds \mathcal{G}_i . That is, for each bid a_{ij} , we compute $\mathbb{E}\left[\frac{a_{ij}-M_j\gamma_i}{a_{ij}} \mid \gamma_i \in \mathcal{G}_i, z_i; \hat{\theta}_\gamma\right]$. Details on this procedure are provided in appendix B.5. Figure 6 plots the distribution of these expected markups across firm-package pairs in green. The expected aggregate markup on awarded bids is 5.2 percent. This point estimate is similar to that of KOW, which estimates an aggregate markup of 4.8 percent on awarded bids.

Finally, we report estimates of costs and markups separately for each awarded bid in table B.6.

²¹We drop the firm's three infeasible dimensions from the proposal distribution, as we describe in appendix B.4.

²²We are able to estimate the joint distribution of firm i 's feasible and infeasible type dimensions because estimation of θ_γ pools information from across firms. However, conditioning on the bounds \mathcal{G}_i is not further informative about the firm's values of those infeasible dimensions.

Figure 6: Distribution of Estimated Bid Markups



Note: Each histogram observation is a bid that a firm i submits on a package j . The blue and orange histograms show the distribution of lower and upper bounds, respectively, on bid markups. The green histogram shows the distribution of mean markups. The blue, orange, and green vertical lines respectively indicate the lower bound, upper bound, and mean markups for the awarded allocation.

5.4.2 Efficient Benchmark

Next, we compare the combinatorial auction to a first-best benchmark by simulating counterfactual outcomes under the Vickrey-Clarke-Groves (VCG) mechanism.²³ The VCG allocation $\tilde{x}_i(c_i, c_{-i}) = \{x_{ij}^*\}_j$ minimizes the total allocation cost, solving

$$\begin{aligned} x^* \in \operatorname{argmin}_x \sum_i \sum_j x_{ij} c_{ij} \\ \text{s.t. } x \in \mathcal{X} \end{aligned} \tag{15}$$

where $x = \{x_{ij}\}_{i,j}$ and $\mathcal{X} \subseteq \prod_i \mathcal{X}_i$ is the same set of allowable allocations imposed in the combinatorial auction (CA). The CA minimizes total payments (the sum of winning bids), while the VCG mechanism minimizes total allocation cost, but both are subject to the allocation constraints imposed in Chile. Under the VCG mechanism, each winning bidder is paid the amount of the (positive) externality generated by its participation in the mechanism: $\tilde{t}_i(c_i, c_{-i}) = \left(\min_x \sum_{i' \neq i} \sum_j x_{i'j} c_{i'j} \right) - \left(\sum_{i' \neq i} \sum_j x_{i'j}^* c_{i'j} \right)$. This is the difference between (i) the minimal allocation cost achievable in firm i 's absence and (ii) the minimal allocation cost when i is included.

For computational tractability, we restrict the set of firm-package pairs that we consider.²⁴ We consider feasible packages for a firm that fall into one of four categories: (i) all packages bid by at least one firm in the observed auction, (ii) all single-TU and two-TU packages, (iii) all packages that contain all TUs in a region, and (iv) all packages containing eight TUs from a single region. These restrictions imply that the cost of this approximate VCG allocation that we simulate is an upper bound on the true minimum cost. However, we expect the approximation error to be small because if complementarities amongst other packages not included in this set were substantial, then at least one firm should have bid the package in the observed auction and the package would be included in set (i).

We estimate economically large efficiency gains in moving from the observed CA to the VCG mechanism. The CA allocation cost is estimated to be about 12.2 percent higher than the (approximate) first-best VCG allocation cost. The VCG mechanism also results

²³In the counterfactual analysis, following KOW, we exclude two firms who submit exceptionally low bids, resulting in high estimated win probabilities and markups. KOW states that “Despite their competitive prices, these firms did not win any units [in the observed auction] and were disqualified from the allocation process because of quality considerations.” We do include these two low-quality firms in the estimation of the bid and type distributions under the assumption that their competitors did not know they would be disqualified at the time of bidding.

²⁴Without any restrictions other than the maximum allowable number of TUs that can be allocated to a firm (eight), there are 100,716,104 possible firm-package pairs.

Table 3: Combinatorial Auction vs. VCG Mechanism

	Mean	SD	p5	p95
<i>Panel A: Combinatorial auction</i>				
Cost	389.3	2.4	385.4	393.1
Producer surplus	21.6	2.4	17.9	25.6
Payment	411.0	-	-	-
<i>Panel B: VCG mechanism</i>				
Cost	341.7	8.3	327.3	353.6
Producer surplus	40.4	7.7	29.0	54.3
Payment	382.1	5.7	372.0	390.1
<i>Panel C: Difference (CA - VCG)</i>				
Cost	47.6	8.1	35.9	61.5
Producer surplus	-18.8	7.6	-32.6	-7.2
Payment	28.9	5.7	20.9	38.9

Note: This table presents summary statistics of simulated CA and VCG outcomes under the estimated type distributions $F_{\gamma_i|z_i}(\cdot | \hat{\theta}^{MH})$, conditioning on the cost bounds \mathcal{G}_i . All values are in pesos per meal.

in higher producer surplus, defined as payments to firms less allocation costs. Although total payments to firms are lower under VCG, this is more than offset by the lower cost of supplying the packages. Table 3 reports summary statistics of simulated allocation costs, producer surpluses, and payments to firms under the observed combinatorial auction and counterfactual VCG mechanism.

To investigate the source of the efficiency gains, we then recompute allocation costs under each auction design under alternative cost economies, holding fixed the simulated VCG allocation draws. We report the results in table 4. In each panel, the first row shows the same baseline allocation cost as in table 3.²⁵

Perhaps the most important source of VCG efficiency gains is the ability to allocate packages with the greatest firm-specific cost complementarities. Shutting down heterogeneity in economies of volume and density barely changes the CA allocation cost but increases the

²⁵Our estimates of the efficiency gains from the VCG allocation are larger than those obtained by KOW, which finds that the combinatorial auction and the VCG auction yield very similar overall costs. A potential reason for the difference is that KOW directly parametrize markups instead of costs. This approach reduces the heterogeneity in markups relative to our model. If we project our estimated markups on the same set of characteristics used in KOW and volume, and recompute the VCG allocation assuming that only packages that a bidder placed a bid on in the combinatorial auction is considered, then the VCG allocation yields a mean cost of 372.2 pesos per meal. Thus, some of the difference between our estimates and those in KOW that arise from the decision to estimate markups versus directly targeting the distribution of costs.

Table 4: CA and VCG Allocation Costs Under Alternative Cost Economies

	Mean	SD	p5	p95
<i>Panel A: Combinatorial auction</i>				
Baseline	389.3	2.4	385.4	393.1
Common economies	390.6	3.4	384.9	396.0
Zero economies	406.1	3.4	400.4	411.6
<i>Panel B: VCG mechanism</i>				
Baseline	341.7	8.3	327.3	353.6
Common economies	371.6	6.0	362.2	381.7
Zero economies	388.1	5.6	379.0	397.8
<i>Panel C: Difference (CA - VCG)</i>				
Baseline	47.6	8.1	35.9	61.5
Common economies	19.0	6.0	9.0	28.5
Zero economies	18.0	5.7	8.6	27.4

Note: In this table, we fix the simulated VCG allocation draws obtained under the estimated type distributions $F_{\gamma_i|z_i}(\cdot | \hat{\theta}^{MH})$. In each row in Panels A and B, we compute summary statistics for the CA or VCG allocation cost under alternative cost economies. The baseline case shows the same allocation cost as in table 3. To compute allocation costs under common cost economies, we set the volume and density cost parameters to their across-firm and across-draw averages. To compute allocation costs under zero cost economies, we set the volume and density cost parameters to be zero. All costs are in pesos per meal.

VCG allocation cost by almost 9 percent. The VCG efficiency gain over the CA is reduced from 12.2 percent to 4.9 percent. These alternative allocation costs are computed by setting each firm’s volume and density cost parameters to the across-firm and across-draw averages of each parameter (see table 4, second row). Eliminating economies of volume and density altogether mechanically increases allocation costs under both mechanisms, but only further reduces the VCG efficiency gain from 4.9 percent to 4.4 percent (see table 4, third row). Thus, the VCG mechanism exploits firm-specific heterogeneity in economies better than the combinatorial auction.

In theory, through package bidding, these firm-specific cost complementarities can also show up in the CA. In practice, the pass-through of these complementarities from package costs to package bids is somewhat limited. This occurs in part because a firm’s bid on a given package j competes with not only its competitors’ bids, but also its own bid on other packages j' . As in the multi-product firm’s pricing problem, the multi-package bidder internalizes the potential for “business stealing” by one of its bids from each of its other bids.

6 Conclusion

A large literature studies the identification of specific econometric models of behavior (see [Matzkin, 2007](#) and the ensuing literature). Ensuring identification is essential in empirical analysis because it is a presumption for the validity of many statistical procedures ([Newey, 1994](#), for example). This paper develops a unified revealed-preference approach for identification in a class of models with a linear payoff structure. The approach is valid in a number of single-agent environments as well as in non-cooperative games; and allows for both continuous and discrete actions, or a combination of the two. We characterize the identified set of models, show how to achieve point identification with excluded shifters, and use our results to suggest an estimation approach.

A number of important examples are a special case of our results. These include single-agent models where revealed preferences are used to identify preferences, for example, in multinomial choice models ([McFadden, 1974, 1981](#)); and reports made to strategy-proof mechanisms such school choice ([Abdulkadiroglu et al., 2017](#)) and second-price auctions. We also cover identification in games when the empirical strategy is based on implications of mutual best-responses in both private information settings (e.g., [Guerre et al., 2000](#)) as well as full-information cases (e.g., [Berry et al., 1995](#)).

These results also apply to cases beyond previously studied settings. As a case in point, our results apply immediately to recent extensions of scoring auctions studied in [Asker and Cantillon \(2008\)](#), including non-linear multi-dimensional scoring auctions as in [Hanazono et al. \(2022\)](#) and scoring rules that combine discrete actions with continuous bids as in [Aspelund and Russo \(2023\)](#). These extensions are useful in a number of settings such as the decision of whether or not to provide add-on service, which is a common feature of a number of auction settings.

Our results also suggest an estimation approach, which we illustrate by revisiting the combinatorial procurement auction for Chilean school lunches studied in [Epstein et al. \(2002\)](#) and KOW. Instead of directly parameterizing bidder markups, our approach targets the distribution of bidder costs and synergies during estimation. This provides an alternative approach in a combinatorial auction setting. We estimate that in the 2003 combinatorial auction, the aggregate markup on awarded bids was 5.2 percent. However, a more efficient allocation was possible. We find that the allocation cost under the status-quo combinatorial auction was 12.2 percent higher than it would have been under the first-best Vickrey-Clarke-Groves mechanism. In theory, package bidding in combinatorial auctions allow bidders to express

across-unit cost complementarities. In practice, in Chile, we find that the pass-through of these complementarities from package costs to package bids is limited. The VCG mechanism’s economically large efficiency gains arise from its ability to take full advantage of across-firm heterogeneity in economies of volume and density.

There are a number of issues that are left for future work. First, a strong restriction in our framework is that payoffs are linear in the outcome space, (x, t) , which consists of an expected allocation and an expected transfer. Linear separability in outcomes and transfer rules out risk aversion. Relaxing this functional form is important. Second, this paper focuses its contributions on identification and largely applies prior methods during estimation. Specifically, we simplified estimation by restricting the dimension of heterogeneity in costs and by parametrizing the cost distribution. Relaxing these assumptions during estimation is also left for future work. Finally, we abstract away from issues such as endogeneity and unobserved heterogeneity, which are useful dimensions in which to extend our approach.

References

- Abdulkadiroglu, Atila, Nikhil Agarwal, and Parag A. Pathak**, “The Welfare Effects of Coordinated School Assignment: Evidence from the NYC High School Match,” *American Economic Review*, 2017, *107*, 3635–89.
- Ackerberg, Daniel A.**, “A new use of importance sampling to reduce computational burden in simulation estimation,” *Quantitative Marketing and Economics*, 12 2009, *7*, 343–376.
- Agarwal, Nikhil and Paulo Somaini**, “Demand Analysis Using Strategic Reports: An Application to a School Choice Mechanism,” *Econometrica*, 2018, *86*, 391–444.
- and —, “Revealed Preference Analysis of School Choice Models,” *Annual Review of Economics*, 8 2020, *12*, 471–501.
- Aliprantis, Charalambos D. and Kim C. Border**, *Infinite dimensional analysis : a hitchhiker’s guide*, 3rd ed., Springer, 2006.
- Allen, Roy and John Rehbeck**, “Identification of Average Demand Models,” *Econometrica*, 2017.
- Asker, John and Estelle Cantillon**, “Properties of scoring auctions,” *The RAND Journal of Economics*, 3 2008, *39*, 69–85.

- Aspelund, Karl and Anna Russo**, “Additionality and Asymmetric Information in Environmental Markets: Evidence from Conservation Auctions,” *Working Paper*, 2023.
- Athey, Susan and Philip A. Haile**, “Identification of Standard Auction Models,” *Econometrica*, 11 2002, 70, 2107–2140.
- **and** —, “Nonparametric Approaches to Auctions,” 1 2007.
- Bajari, Patrick and C. Lanier Benkard**, “Demand Estimation with Heterogeneous Consumers and Unobserved Product Characteristics: A Hedonic Approach,” *Journal of Political Economy*, 12 2005, 113, 1239–1276.
- Berry, Steven and Philip Haile**, “Identification in Differentiated Products Markets,” *Annual Review of Economics*, 10 2016, 8, 27–52.
- Berry, Steven T. and Philip A. Haile**, “Identification in Differentiated Products Markets Using Market Level Data,” *Econometrica*, 2014, 82, 1749–1798.
- **and** —, “Identification of Nonparametric Simultaneous Equations Models With a Residual Index Structure,” *Econometrica*, 2018, 86, 289–315.
- , **James Levinsohn, and Ariel Pakes**, “Automobile Prices in Market Equilibrium,” *Econometrica*, 1995, 63, 841 – 890.
- Cantillon, Estelle and Martin Pesendorfer**, “Combination Bidding in Multi-Unit Auctions,” 2007.
- Ciliberto, Federico and Elie Tamer**, “Market Structure and Multiple Equilibria in Airline Markets,” *Econometrica*, 2009, 77, 1791–1828.
- Dieudonné, Jean**, *Treatise on Analysis*, 1 ed., Vol. II, Academic Press, Inc., 1976.
- Epstein, Rafael, Lysette Henríquez, Jaime Catalán, Gabriel Y. Weintraub, and Cristián Martínez**, “A Combinational Auction Improves School Meals in Chile,” *Interfaces*, 12 2002, 32, 1–14.
- Gentry, Matthew L., Tatiana Komarova, and Pasquale Schiraldi**, “Simultaneous First-Price Auctions with Preferences Over Combinations: Identification, Estimation and Application,” *SSRN Electronic Journal*, 10 2014.
- Gorman, W. M.**, “A Possible Procedure for Analysing Quality Differentials in the Egg Market,” *The Review of Economic Studies*, 10 1980, 47, 843–856.

- Guerre, Emmanuel, Isabelle Perrigne, and Quang Vuong**, “Optimal Nonparametric Estimation of First-Price Auctions,” *Econometrica*, 2000, *68*, 525–574.
- Hanazono, Makoto, Yohsuke Hirose, Jun Nakabayashi, and Masanori Tsuruoka**, “Theory and Estimation for Scoring Auctions,” 11 2022.
- Kim, Sang Won, Marcelo Olivares, and Gabriel Y. Weintraub**, “Measuring the Performance of Large-Scale Combinatorial Auctions: A Structural Estimation Approach,” *Management Science*, 5 2014, *60*, 1180–1201.
- Lancaster, Kelvin J.**, “A New Approach to Consumer Theory on JSTOR,” *Journal of Political Economy*, 4 1966, *74*, 132–157.
- Larsen, Bradley and Anthony Lee Zhang**, “A Mechanism Design Approach to Identification and Estimation,” 2018.
- Lewbel, Arthur**, “Semiparametric qualitative response model estimation with unknown heteroscedasticity or instrumental variables,” *Journal of Econometrics*, 7 2000, *97*, 145–177.
- Matzkin, Rosa L.**, “Nonparametric identification,” 1 2007.
- McFadden, Daniel**, “Conditional Logit Analysis of Qualitative Choice Behavior,” 1973, pp. 105 – 142.
- , “Conditional logit analysis of qualitative choice behavior,” *Frontiers in econometrics*, 1974, pp. 105–142.
- , “Econometric models of probabilistic choice,” *Structural analysis of discrete data with econometric applications*, 1981.
- Milgrom, Paul and Ilya Segal**, “Envelope Theorems for Arbitrary Choice Sets,” *Econometrica*, 2002, *70*, 583 – 601.
- Newey, Whitney K.**, “The Asymptotic Variance of Semiparametric Estimators,” *Econometrica*, 11 1994, *62*, 1349.
- OECD**, “Main Economic Indicators - complete database,” 2016.
- Pakes, A.**, “Alternative Models for Moment Inequalities,” *Econometrica*, 2010, *78*, 1783–1822.

Rockafellar, R. Tyrrell, *Convex Analysis*, Princeton University Press, 1970.

Rosen, Sherwin, “Hedonic Prices and Implicit Markets: Product Differentiation in Pure Competition on JSTOR,” *Journal of Political Economy*, 1 1974, 82, 34–55.

Shabat, Boris Vladimirovich, *Introduction to Complex Analysis, Part 2*, American Mathematical Society, 1992.

Tebaldi, Pietro, Alexander Torgovitsky, and Hanbin Yang, “Nonparametric Estimates of Demand in the California Health Insurance Exchange,” 5 2019.

Train, Kenneth E., “Discrete Choice Methods with Simulation,” *Cambridge Books*, 2009.

Xiao, Mo and Zhe Yuan, “License Complementarity and Package Bidding: The U.S Spectrum Auctions,” 2020.

Appendix

A Lemmata

Lemma 2. *Let $t : \mathbb{R}^J \rightarrow \mathbb{R} \cup \{\infty\}$ be a proper closed convex function supported over a convex compact set \mathcal{X} that has strictly non empty interior in \mathbb{R}^J . Then there exist $B \subset \mathcal{X}$ such that $\partial t(B) = \{v \in \mathbb{R}^J : v \in \partial t(x), x \in B\}$ has a non empty interior in \mathbb{R}^J and is contained in a translated simplicial cone.*

Proof. The conjugate function t^* is a proper closed convex function [Rockafellar \(1970, Theorem 12.2\)](#). Because t is proper and lower semi-continuous [Rockafellar \(1970, page 52\)](#), and \mathcal{X} is compact, for every $v \in \mathbb{R}^J$, $t^*(v) = \sup_{x \in \mathcal{X}} v \cdot x - t(x)$ is attained for some $x \in \mathcal{X}$. and $\partial t^*(v) \subset \mathcal{X}$. Therefore, the support of t^* is \mathbb{R}^J . The set D where the function f^* is differentiable is a dense subset of \mathbb{R}^J , and the gradient mapping $\nabla t^* : v \rightarrow \nabla t^*(v)$ is continuous on D [Rockafellar \(1970, Theorem 25.5\)](#). Take any $v_0 \in D$ and let $x_0 = \nabla t^*(v_0) = \partial t^*(v_0) \in \mathcal{X}$. Let $S = \{x_1, x_2, \dots, x_J\} \subset \mathcal{X}$ be such that the convex hull of $S \cup \{x_0\}$ has strictly positive Lebesgue measure in \mathbb{R}^J . Such set S exists because \mathcal{X} has strictly positive Lebesgue measure in \mathbb{R}^J . Let $\bar{x} \in \mathbb{R}^J$ be such that x_0 is in the interior of the convex hull of $S \cup \{\bar{x}\}$. Such an \bar{x} exists although it may not be an element in \mathcal{X} . Let $\bar{B}(x_0)$ be neighborhood of x_0 in the interior of the convex hull of $S \cup \{\bar{x}\}$. By the continuity of the gradient mapping ∇t^* , there exist a neighborhood, $\bar{V}(v_0) \subset \mathbb{R}^J$ such that $B = \{x \in \mathbb{R}^J : x = \nabla t^*(v), v \in \bar{V}(v_0)\} \subset \bar{B}(x_0)$. Because for all $x \in B$, $x \in \partial t^*(v) \subset \mathcal{X}$ for some $v \in \bar{V}(v_0)$, it follows that $B \subset \mathcal{X}$. Moreover, if $v \in \bar{V}(v_0)$, then $x \in \partial t^*(v) \subset B$ [Rockafellar \(1970, Theorem 25.6\)](#). Therefore, $v \in \partial t(x) \subset \partial t(B)$ [Rockafellar \(1970, Theorem 23.5\)](#). Thus, $\partial t(B) \supset \bar{V}(v_0)$ has strictly positive Lebesgue measure in \mathbb{R}^J . \square

The conjugate function t^* is a proper closed convex function [Rockafellar \(1970, Theorem 12.2\)](#). Because t is proper and lower semi-continuous [Rockafellar \(1970, page 52\)](#), and \mathcal{X} is compact, for every $v \in \mathbb{R}^J$, $t^*(v) = \sup_{x \in \mathcal{X}} v \cdot x - t(x)$ is attained for some $x \in \mathcal{X}$. and $\partial t^*(v) \subset \mathcal{X}$. Therefore, the support of t^* is \mathbb{R}^J . The set D where the function f^* is differentiable is a dense subset of \mathbb{R}^J , and the gradient mapping $\nabla t^* : v \rightarrow \nabla t^*(v)$ is continuous on D [Rockafellar \(1970, Theorem 25.5\)](#). Take any $v_0 \in D$ and let $x_0 = \nabla t^*(v_0) = \partial t^*(v_0) \in \mathcal{X}$. Let $S = \{x_1, x_2, \dots, x_J\} \subset \mathcal{X}$ be such that the convex hull of $S \cup \{x_0\}$ has strictly positive Lebesgue measure in \mathbb{R}^J . Such set S exists because \mathcal{X} has strictly positive Lebesgue measure in \mathbb{R}^J . Let $\bar{x} \in \mathbb{R}^J$ be such that x_0 is in the interior of the convex hull of $S \cup \{\bar{x}\}$. Such

an \bar{x} exists although it may not be an element in \mathcal{X} . Let $\bar{B}(x_0)$ be neighborhood of x_0 in the interior of the convex hull of $S \cup \{\bar{x}\}$. By the continuity of the gradient mapping ∇t^* , there exist a neighborhood, $\bar{V}(v_0) \subset \mathbb{R}^J$ such that $B = \{x \in \mathbb{R}^J : x = \nabla t^*(v), v \in \bar{V}(v_0)\} \subset \bar{B}(x_0)$. Because for all $x \in B$, $x \in \partial t^*(v) \subset \mathcal{X}$ for some $v \in \bar{V}(v_0)$, it follows that $B \subset \mathcal{X}$. Moreover, if $v \in \bar{V}(v_0)$, then $x \in \partial t^*(v) \subset B$ [Rockafellar \(1970, Theorem 25.6\)](#). Therefore, $v \in \partial t(x) \subset \partial t(B)$ [Rockafellar \(1970, Theorem 23.5\)](#). Thus, $\partial t(B) \supset \bar{V}(v_0)$ has strictly positive Lebesgue measure in \mathbb{R}^J .

Now, we show that B is contained in a translated simplicial cone. Consider the choice set $\mathcal{Y} = S \cup \{\bar{x}\}$ with transfers $f_{\mathcal{Y}}(x) = t(x)$ for $x \in S$ and $f_{\mathcal{Y}}(\bar{x})$ sufficiently low so that all outcomes in $\bar{B}(x_0)$ with transfers $t(\cdot)$ would be dominated by the outcomes in \mathcal{Y} . Let $f(\cdot)$ be the convex hull of the function $f_{\mathcal{Y}}(\cdot)$. The function $f(\cdot)$ is the largest convex function that lies below $f_{\mathcal{Y}}(\cdot)$ and coincides with $f_{\mathcal{Y}}$ on the set \mathcal{Y} . The subdifferential $\partial f(\bar{x})$ is a translated simplicial cone because the set \mathcal{Y} includes $J+1$ affinely independent points in \mathbb{R}^J . If $v \in \partial t(B)$, then an agent with value v weakly prefers some $x \in B$ to any $x' \in S$. However, by construction, dominance implies that some element of $S \cup \{\bar{x}\}$ is strictly preferred to x . Thus, the agent strictly prefers \bar{x} to x and, by transitivity, to all $x' \in S$. Therefore, $\partial t(B) \subset \partial f(\bar{x})$ which is a translated simplicial cone.

Lemma 3. *Let C be a subset of $\mathcal{X} \subseteq \mathbb{R}^J$ with strictly positive Lebesgue measure. Assume that $C \subseteq \bar{C} + \{x\}$, where \bar{C} is a simplicial cone and $+$ denotes Minkowski summation. Then, there exists $\lambda \in \text{int } \mathcal{N}(\bar{C})$, such that*

$$\hat{\chi}_{C,\lambda}(\xi) = \int_C \exp(-2\pi v \cdot (i\xi + \lambda)) dv$$

is not zero on any open set $\Xi \subseteq \mathbb{R}^J$. Moreover, $\hat{\chi}_{C,\tilde{\lambda}}(\xi)$ is not zero on any open set $\Xi \subseteq \mathbb{R}^J$ for any $\tilde{\lambda} = \alpha\lambda$ for $\alpha \in (0, 1)$.

Proof. We first introduce some notation. First, define the matrix $A_{\bar{C}} = \begin{bmatrix} a_1 \\ \vdots \\ a_J \end{bmatrix}$ so that $v \in \bar{C}$

if and only if $v = A_{\bar{C}}u$ for some $u \geq 0$. By definition, $A_{\bar{C}}$ is invertible because \bar{C} is simplicial. It is without loss of generality to assume that $|\det(A_{\bar{C}})| = 1$. Second, let $\mathcal{N}(\bar{C})$ be the dual cone to \bar{C} . Note that $\text{int } \mathcal{N}(\bar{C})$ is non-empty because \bar{C} is simplicial. Fix a $\lambda \in \text{int } \mathcal{N}(\bar{C})$ for the remainder of this proof.

Towards a contradiction, assume that $\hat{\chi}_{C,\lambda}(\xi)$ is zero on an open set $\Xi \subseteq \mathbb{R}^J$. We will show below that $\hat{\chi}_{C,\lambda}(\xi)$ when viewed as a function on $\Xi_\lambda = \{\xi \in \mathbb{C}^J : \xi = y + iz, y \in \Xi\}$

$\mathbb{R}^J, \frac{\lambda}{2} - z \in \text{int } \mathcal{N}(\bar{C})\}$, which contains \mathbb{R}^J , is holomorphic. Under this hypothesis, Theorem 5 in [Shabat \(1992\)](#) implies that $\hat{\chi}_{C,\lambda}(\xi) = 0$ for all $\xi \in \mathbb{R}^J$. However, this contradicts the fact that $\hat{\chi}_{C,\lambda}(\xi)$ is, up to scale, the characteristic function of a random variable with density $1 \{v \in C\} \exp(-2\pi \langle v, \lambda \rangle)$. This density, and therefore $\hat{\chi}_{C,\lambda}(\xi)$, is non-zero because C has strictly positive Lebesgue measure.

The second part follows immediately because if $\lambda \in \text{int } \mathcal{N}(\bar{C})$ then so is $\tilde{\lambda} = \alpha\lambda$ for any $\alpha \in (0, 1)$.

Hence, it only remains to show that $\hat{\chi}_{C,\lambda}(\xi)$ is holomorphic in Ξ_λ . To do this, we first use the differentiation under the integral sign theorem for complex variables (Theorem 13.8.6(iii) in [Dieudonné, 1976](#)) to show that $\frac{\partial \hat{\chi}_{C,\lambda}(\xi)}{\partial \xi_k}$ exists on Ξ_λ and is equal to

$$\frac{\partial \hat{\chi}_{C,\lambda}(\xi)}{\partial \xi_k} = \int 1 \{v \in C\} i v_k \exp(-2\pi v \cdot (i\xi + \lambda)) dv.$$

Fix $\xi_{-k} = (\xi_1, \dots, \xi_{k-1}, \xi_{k+1}, \dots, \xi_J)$ and define the function $f_C(v, \xi_k; \xi_{-k}) = 1 \{v \in C\} \exp(-2\pi v \cdot (i\xi + \lambda))$. To apply the result, we need to show that $f_C(v, \xi_k; \xi_{-k})$ is (i) analytic in ξ_k for almost all v , (ii) measurable in v for each ξ , and (iii) there exists an integrable function $g(v)$ such that for almost all ξ , $|f_C(v, \xi_k; \xi_{-k})| \leq g(v)$. Requirement (i) follows from the definition of $f_C(\cdot)$. Measurability is immediate given the definition of $f_C(v, \xi_k; \xi_{-k})$.

To show (iii) we will use the fact that $|f_{\bar{C}+\{x\}}(v, \xi_k; \xi_{-k})| \geq |f_C(v, \xi_k; \xi_{-k})|$ for all v and show that $|f_{\bar{C}+\{x\}}(v)| \leq g(v) = 1 \{v - x \in \bar{C}\} \left| \exp\left(-2\pi v \cdot \frac{\lambda}{2}\right) \right|$ for all $\xi \in \Xi_\lambda$, where $g(v)$ is integrable. To this end, we first bound $|f_{\bar{C}+\{x\}}(v)|$ as follows:

$$\begin{aligned} |f_{\bar{C}+\{x\}}(v)| &= \left| 1 \{v - x \in \bar{C}\} \exp(-2\pi v \cdot (i\xi + \lambda)) \right| \\ &= 1 \{v - x \in \bar{C}\} \left| \exp(-2\pi v \cdot (i(y + iz) + \lambda)) \right| \\ &= 1 \{v - x \in \bar{C}\} \left| \exp\left(-2\pi v \cdot \left(\frac{\lambda}{2} + \left(\frac{\lambda}{2} - z\right)\right)\right) \right| \\ &\leq 1 \{v - x \in \bar{C}\} \left| \exp\left(-2\pi v \cdot \frac{\lambda}{2}\right) \right| \equiv g(v). \end{aligned}$$

The first equality uses the definition $\xi = y + iz$, The second equality follows the fact that

$$\begin{aligned} |\exp(-2\pi v \cdot (i(y + iz) + \lambda))| &= |\exp(-2\pi v \cdot iy)| |\exp(-2\pi v \cdot (\lambda - z))| \\ &= |\exp(-2\pi v \cdot (\lambda - z))|. \end{aligned}$$

The inequality uses the fact that $\frac{\lambda}{2} - z \in \text{int } \mathcal{N}(\bar{C})$, implying that $v \cdot \left(\frac{\lambda}{2} - z\right) > 0$. We now show that $g(v)$ is integrable.

$$\begin{aligned}
\int g(v) dv &= \int 1 \{v - x \in \bar{C}\} \left| \exp \left(-2\pi v \cdot \frac{\lambda}{2} \right) \right| dv \\
&= \int 1 \{A_{\bar{C}}^{-1}(v - x) \geq 0\} \left| \exp \left(-2\pi v \cdot \frac{\lambda}{2} \right) \right| dv \\
&= \int_{\mathbb{R}_+^J} \left| \exp \left(-2\pi (A_{\bar{C}} u + x) \cdot \frac{\lambda}{2} \right) \right| du \\
&= \exp \left(-2\pi x \cdot \frac{\lambda}{2} \right) \int_{\mathbb{R}_+^J} \exp \left(-2\pi A_{\bar{C}} u \cdot \frac{\lambda}{2} \right) du.
\end{aligned}$$

The first equality uses the definition of $A_{\bar{C}}$. The second substitutes $u = A_{\bar{C}}^{-1}(v - x)$, and uses the fact that $|\det(A_{\bar{C}})| = 1$ and $u \geq 0$. The third rewrites the integrands as the product of two terms and pulls out the constant $\left| \exp \left(-2\pi x \cdot \frac{\lambda}{2} \right) \right|$ from the integral sign.

Since $\exp \left(-2\pi x \cdot \frac{\lambda}{2} \right)$ is finite, we need to show that $\int_{\mathbb{R}_+^J} \exp \left(-2\pi A_{\bar{C}} u \cdot \frac{\lambda}{2} \right) du$ is finite. Observe that $\frac{\lambda}{2} \in \text{int } \mathcal{N}(\bar{C})$. Re-write the integral as

$$\begin{aligned}
&\int_{\mathbb{R}_+^J} \exp \left(-2\pi A_{\bar{C}} u \cdot \frac{\lambda}{2} \right) du \\
&= \int_{\mathbb{R}_+^J} \exp \left(-2\pi \|u\| A_{\bar{C}} \frac{u}{\|u\|} \cdot \frac{\lambda}{2} \right) du \\
&= \int_{\mathbb{S}_+^{J-1}} \int_0^\infty r^{J-1} \exp \left(-2\pi r A_{\bar{C}} u' \cdot \frac{\lambda}{2} \right) dr d\sigma(u') \\
&= \int_{\mathbb{S}_+^{J-1}} \frac{1}{(2\pi A_{\bar{C}} u' \cdot \frac{\lambda}{2})^J} \int_0^\infty s^{J-1} \exp(-s) ds d\sigma(u') \\
&= \int_{\mathbb{S}_+^{J-1}} \frac{1}{(2\pi A_{\bar{C}} u' \cdot \frac{\lambda}{2})^J} \Gamma(J) d\sigma(u'),
\end{aligned}$$

where $\sigma(u')$ is the proper surface measure of the sphere \mathbb{S}^{J-1} , $\mathbb{S}_+^{J-1} = \mathbb{S}^{J-1} \cap \mathbb{R}_+^J$, $u' = \frac{u}{\|u\|}$, and $\Gamma(\cdot)$ is the Gamma function. The first equality is trivial, the second from a change of variables $u' = \frac{u}{\|u\|}$ and $r = \|u\|$, the third from a change of variables $s = -2\pi r A_{\bar{C}} u' \cdot \frac{\lambda}{2}$ and the last from the definition of the Gamma function.

Because $\frac{\lambda}{2} \in \text{int } \mathcal{N}(\bar{C})$, we have that $(A_{\bar{C}} u' \cdot \frac{\lambda}{2})$ is strictly positive, with a strictly positive

minimum $\kappa > 0$. Hence, we have that

$$\int_{\mathbb{S}_+^{J-1}} \frac{1}{(2\pi A_{\bar{C}} u' \cdot \frac{\lambda}{2})^J} \Gamma(J) d\sigma(u') < (2\pi\kappa)^{-J} \Gamma(J) \int_{\mathbb{S}_+^{J-1}} 1 d\sigma(u').$$

Since $\sigma(\mathbb{S}^{J-1})$ is finite, we have that the right hand side is finite.

Finally, Osgood's Lemma implies that $\hat{\chi}_{C,\lambda}(\xi)$ is holomorphic if it is continuous in ξ . For any $h \in \mathbb{C}^J$,

$$|\hat{\chi}_{C,\lambda}(\xi + h) - \hat{\chi}_{C,\lambda}(\xi)| \leq \int 1\{v \in C\} |\exp(-2\pi v \cdot (i(\xi + h) + \lambda)) - \exp(-2\pi v \cdot (i\xi + \lambda))| dv.$$

Since $\xi \in \Xi_\lambda$, there exists $\varepsilon > 0$ such that for $h \in \mathbb{C}^J$ with $\|h\| < \varepsilon$, $\xi + h \in \Xi_\lambda$. As both $\xi + h$ and ξ are in Ξ_λ , the integrand is dominated by $2g(v)$, which is integrable. And $1\{v \in C\} |\exp(-2\pi v \cdot (i(\xi + h) + \lambda)) - \exp(-2\pi v \cdot (i\xi + \lambda))| = \exp(-2\pi v \cdot \lambda) |\exp(-2\pi v \cdot ih) - 1| \rightarrow 0$, as $h \rightarrow 0$. Therefore, by the dominated convergence theorem, $|\hat{\chi}_{C,\lambda}(\xi + h) - \hat{\chi}_{C,\lambda}(\xi)| \rightarrow 0$ as $h \rightarrow 0$. \square

B Empirical Appendix

B.1 Combinatorial Auction Mechanism

In this section, we present the full winner determination problem, as presented in KOW's appendix G. The decision variables $x_{ij} \in \{0, 1\}$ indicate whether firm i is allocated package j . The combinatorial auction minimizes total payments (the sum of winning bids) by solving the integer program

$$\begin{aligned}
& \min_x \sum_i \sum_j x_{ij} a_{ij} \\
& \text{s.t.} \quad \sum_i \sum_{j:k \in j} x_{ij} \geq 1 \quad \text{for all } k \\
& \quad \sum_j x_{ij} \leq 1 \quad \text{for all } i \\
& \quad \sum_j x_{ij} q_{ij} \leq \bar{q}_i \quad \text{for all } i \\
& \quad \sum_j x_{ij} |j| \leq \bar{n}_i \quad \text{for all } i \\
& \quad \underline{I}_r \leq \sum_i \sum_{j:j \cap r \neq \emptyset} x_{ij} \leq \bar{I}_r \quad \text{for all } r \\
& \quad \sum_i \sum_j x_{ij} \geq \underline{I}
\end{aligned}$$

In words, the constraints are:

1. Each TU k is assigned;
2. Each firm i wins at most one package;
3. Each firm i wins at most \bar{q}_i total meals;
4. Each firm i wins at most \bar{n}_i total TUs;
5. Each region r is served by between \underline{I}_r and \bar{I}_r firms; and
6. At least \underline{I} total firms are included in the allocation

In KOW, the authors write that the second constraint isn't explicitly imposed in the actual auctions, though it turns out that this constraint is not binding in most years. We follow KOW in including this second constraint. We never observe \underline{I} , the minimum number of firms that must win in a given auction, needed to check the final constraint; we drop this final constraint in our application.

Table B.1: Estimates of Bid Adjustment Function Parameters β^{volume} and β^{density}

	Small firms	Large firms
Volume 1	-12.899 (0.929)	-16.158 (1.085)
Volume 2	-17.752 (0.934)	-22.742 (1.077)
Volume 3	-20.429 (0.937)	-27.144 (1.076)
Volume 4	-25.683 (1.234)	-29.265 (1.077)
Density 1	3.919 (0.125)	0.797 (0.076)
Density 2	5.629 (0.155)	1.094 (0.091)

Note: Standard errors are in parentheses.

B.2 Bid Distribution

B.2.1 Bid Adjustment Functions

The bid adjustment functions $h_a^{\text{volume}}(\cdot; \beta_{\text{size}_i}^{\text{volume}})$ and $h_a^{\text{density}}(\cdot; \beta_{\text{size}_i}^{\text{density}})$ are step functions in the volume q_j and density d_j , respectively, of the package j . The volume adjustment function is constant in each of nine equally spaced intervals $\{[0, 3], \dots, [24, 27]\}$ that cover the range of package volumes observed in the data. The density adjustment function is constant in each of the intervals $\{[0, 0.5), [0.5, 1), [1, 1]\}$, with the third bin capturing the mass of packages whose constituent TUs lie entirely within one region.²⁶ We normalize bid adjustments in the smallest volume bin and largest density bin to zero.

B.2.2 Parameter Estimates


In tables B.1, B.2, and B.3, we present estimates $\hat{\theta}_a$ of the bid distribution parameters. In figure B.1, we compare the observed distribution of per-meal prices bid to our estimated distribution.

²⁶Package volume is measured in millions of meals per year. We define package density in footnote 11; it takes values in the unit interval.

Table B.2: Estimates of Standard Deviations σ_ϵ of Idiosyncratic Bid Shocks

Package size (# TUs)	Standard devia- tion
1	13.757
2	4.870
3	3.901
4	3.211
5	2.523
6	2.054
7	1.319
8	1.502

Figure B.1: Observed and Estimated Bid Distributions



Note: The blue histogram shows prices bid by all firms on all packages on which they submitted bids in the data; an observation is a firm-package pair. To construct the orange histogram, for each firm-package pair, we simulate 1000 draws from the estimated bid distribution; each observation is a firm-package-draw triple. Each histogram is normalized to represent a discrete probability distribution.

Table B.3: Estimates of Means β^{TU} and Standard Deviations σ_ω of TU Base Prices

	β	σ_ω
TU 401	500.949	14.839
TU 402	440.899	12.154
TU 403	515.466	13.684
TU 404	606.711	33.822
TU 405	580.172	33.026
TU 501	501.379	30.620
TU 502	421.620	19.436
TU 503	430.904	17.976
TU 504	417.353	21.962
TU 505	433.111	11.867
TU 506	445.616	13.667
TU 507	450.197	12.639
TU 508	404.258	14.435
TU 510	416.788	9.913
TU 511	433.948	29.586
TU 901	510.563	25.836
TU 902	512.277	24.509
TU 903	585.829	36.108
TU 904	443.816	18.065
TU 905	486.187	22.731
TU 906	474.471	19.505
TU 907	499.643	24.880
TU 908	528.013	24.865
TU 909	581.423	38.627
TU 1201	603.123	48.260
TU 1331	415.935	13.356
TU 1332	378.521	11.529
TU 1333	356.387	11.532
TU 1334	404.167	11.000
TU 1335	392.525	21.249
TU 1336	404.571	12.959
TU 1339	444.407	20.426
Incumbency shifter	-17.621	

Note: The variance-covariance matrix Σ_ω is unrestricted, but for clarity only the (square roots of the) diagonal elements are reported here.

B.3 Type Bounds

As part of estimating the type bounds $\tilde{\mathcal{G}}_i$, we rule out downward deviations to bid vectors a'_i in which firm i decreases its bid on a single package j by 50 pesos per meal. The set of packages $\tilde{\mathcal{J}}_i$ for which we consider these downward deviations is chosen to generate independent variation in each of the type dimensions. Specifically, for each of the type dimensions indexed by l , we start by choosing a set of packages $\tilde{\mathcal{J}}_{il}$ to be maximally informative about that dimension:

- If l is a volume or density economy, then $\tilde{\mathcal{J}}_{il}$ is the set of 10 packages with the highest estimated win probabilities for firm i under its observed bids.
- If l is a TU cost, then $\tilde{\mathcal{J}}_{il}$ contains only the single-unit package with that TU. If we do not observe firm i bidding on the single-unit package for that TU, then the sole element of $\tilde{\mathcal{J}}_{il}$ is the next-smallest package i bids on, with ties broken in descending order of i 's estimated probability of winning that package.

Then, we take the union of these dimension-specific sets $\tilde{\mathcal{J}}_{il}$ and further restrict to packages that (i) firm i bids on in the data, (ii) firm i is allowed to win under the meal and TU constraints described in appendix B.1, and (iii) whose costs have a nonzero coefficient on dimension l .

Note that not all firms are allowed to win packages involving all type dimensions. For example, some firms are restricted in the size of packages they can win (where size is measured both in number of meals and number of TUs). If firm i is not allowed to win any packages falling in the highest volume bin, then revealed preference can tell us nothing about the element of γ_i^{volume} corresponding to that volume bin.

B.4 Type Distribution

B.4.1 Importance Sampling

In principle, it suffices to estimate the likelihood (13) at each candidate θ_γ by counting how many draws from $F_{\Gamma_i|z_i}(\cdot | \theta_\gamma)$ lie in the set \mathcal{G}_i , though a precise estimate of a small likelihood – if $F_{\Gamma_i|z_i}(\cdot | \theta_\gamma)$ puts little mass on \mathcal{G}_i – may require a large number of draws. We would ideally like to sample γ_i from $F_{\Gamma_i|z_i}(\cdot | \theta_\gamma)$ conditional on γ being in \mathcal{G}_i , but doing so in closed

form is possible only in one dimension. Instead, we use importance sampling, taking draws from a proposal distribution G_i and reweighting them by their likelihood ratios:

$$L(\theta_\gamma \mid \mathcal{G}_i) = \int 1\{x \in \mathcal{G}_i\} \frac{f_{\Gamma_i|z_i}(x; \theta_\gamma)}{g_i(x)} dG_i(x) \quad (14)$$

We furthermore use draws from G_i conditional on the type bounds \mathcal{G}_i ; denote this conditional distribution by \tilde{G}_i . Inside the set \mathcal{G}_i , the unconditional and conditional distributions differ by the total mass that G_i puts on \mathcal{G}_i :

$$d\tilde{G}_i(x) = \frac{1\{x \in \mathcal{G}_i\}}{\int 1\{y \in \mathcal{G}_i\} dG_i(y)} dG_i(x) \quad (16)$$

In particular, when $x \in \mathcal{G}_i$, then $dG_i(x) = [\int 1\{y \in \mathcal{G}_i\} dG_i(y)] d\tilde{G}_i(x)$. As a result, the likelihood in equation (14) becomes

$$\begin{aligned} L(\theta_\gamma \mid \mathcal{G}_i) &= \int 1\{x \in \mathcal{G}_i\} \frac{f_{\Gamma_i|z_i}(x; \theta_\gamma)}{g_i(x)} \left[\int 1\{y \in \mathcal{G}_i\} dG_i(y) \right] d\tilde{G}_i(x) \\ &= \left[\int 1\{y \in \mathcal{G}_i\} dG_i(y) \right] \times \int \frac{f_{\Gamma_i|z_i}(x; \theta_\gamma)}{g_i(x)} d\tilde{G}_i(x) \\ &=: \left[\int 1\{y \in \mathcal{G}_i\} dG_i(y) \right] \times \tilde{L}(\theta_\gamma \mid \tilde{\Gamma}_i) \end{aligned}$$

where the second equality uses the fact that, by definition, $1\{x \in \mathcal{G}_i\} \equiv 1$ for x drawn from \tilde{G}_i . Call $\tilde{L}(\theta_\gamma \mid \mathcal{G}_i)$ the **quasi-likelihood**, which differs from the true likelihood by the same factor $\int 1\{y \in \mathcal{G}_i\} g_i(y) dy$ as in equation (16). Since this factor does not depend on θ_γ , it suffices to use the quasi-likelihood in Metropolis-Hastings rather than the true likelihood. The quasi-likelihood is easy to compute, as it doesn't require integrating out the mass that the unconditional proposal distribution puts on \mathcal{G}_i . We approximate the quasi-likelihood using $S_\gamma^{\text{prop}} = 1,000$ draws $\gamma_i^{(s)}$ from \tilde{G}_i , the version of the proposal distribution that conditions on the type bounds:

$$\tilde{L}(\theta_\gamma \mid \tilde{\Gamma}_i) \approx \frac{1}{S_\gamma^{\text{prop}}} \sum_{s=1}^{S_\gamma^{\text{prop}}} \frac{f_{\Gamma_i|z_i}(\gamma_i^{(s)}; \theta_\gamma)}{g_i(\gamma_i^{(s)})}$$

Appendix B.4.2 describes how we generate draws from the conditional proposal distribution.

Finally, this discussion has so far assumed that firm i is allowed to win at least one package involving each type dimension. If there are in fact some dimensions that are infeasible for i , then \mathcal{G}_i must always be completely unbounded in those dimensions, as no bids by the

firm can ever be informative about its values for those dimensions. When evaluating the (quasi-)likelihood, we therefore drop infeasible dimensions from both the type bound \mathcal{G}_i and the proposal distribution G_i , and we integrate out over infeasible dimensions in $F_{\Gamma_i|z_i}$. (This is easy because G_i is a product distribution and $F_{\Gamma_i|z_i}$ is multivariate normal.)

B.4.2 Proposal Distribution

Constructing the Unconditional Distribution The advantage of importance sampling is that fewer draws are required to precisely estimate the likelihood; the challenge is choosing the proposal distribution well. Its support must contain \mathcal{G}_i , and it ideally concentrates most of its mass there. Its density should be easy to evaluate. The variance of the resulting likelihood ratios should be minimized, so that the effective sample size is maximized. The difficulty of designing the right proposal distribution increases with the dimension of the random vector. Let G_i be a (firm-specific) product distribution on the L -dimensional box containing \mathcal{G}_i . We use a product of uniform and exponential distributions, depending on whether the given dimension is bounded. For each dimension l , define

$$\begin{aligned}\varpi_{il} &= \inf \left\{ \gamma_{il} \in \mathbb{R} \cup \{-\infty\} : (\gamma_{il}, \gamma_{i,-l}) \in \mathcal{G}_i \text{ for some } \gamma_{i,-l} \in \mathbb{R}^{L-1} \right\} \\ \bar{\varpi}_{il} &= \sup \left\{ \gamma_{il} \in \mathbb{R} \cup \{+\infty\} : (\gamma_{il}, \gamma_{i,-l}) \in \mathcal{G}_i \text{ for some } \gamma_{i,-l} \in \mathbb{R}^{L-1} \right\} \\ \chi_{il} &= \begin{cases} -100, & l \text{ is a volume or density economy} \\ 300, & l \text{ is a TU cost} \end{cases} \\ \bar{\chi}_{il} &= \begin{cases} 100, & l \text{ is a volume or density economy} \\ 600, & l \text{ is a TU} \end{cases}\end{aligned}$$

Then we define G_{il} , the proposal distribution for dimension l , as follows:

- If $\varpi_{il} > \chi_{il}$ and $\bar{\varpi}_{il} < \bar{\chi}_{il}$, then G_{il} is uniform on $[\varpi_{il}, \bar{\varpi}_{il}]$.
- If $\varpi_{il} > \chi_{il}$ but not $\bar{\varpi}_{il} < \bar{\chi}_{il}$, then we take G_{il} to be unbounded from above. Let G_{il} be the shifted and re-scaled exponential distribution with minimum ϖ_{il} and 75th percentile $\bar{\chi}_{il}$.
- If $\bar{\varpi}_{il} < \bar{\chi}_{il}$ but not $\varpi_{il} > \chi_{il}$, then we take G_{il} to be unbounded from below. Let G_{il} be the shifted, rescaled, and flipped exponential distribution with maximum $\bar{\varpi}_{il}$ and 25th percentile χ_{il} .

- If neither $\varrho_{il} > \chi_{il}$ nor $\bar{\gamma}_{il} < \bar{\chi}_{il}$, then we take G_{il} to be unbounded in both directions. Let G_{il} be the shifted and re-scaled Laplace (double exponential) distribution with 25th percentile χ_{il} and 75th percentile $\bar{\chi}_{il}$.

Ideally, we would like to sample uniformly from the smallest box in \mathbb{R}^L containing \mathcal{G}_i , which would amount to sampling uniformly and independently from each dimension. This is certainly not possible in dimension l if \mathcal{G}_i is unbounded in that dimension—that is, if the **outer bounds** ϱ_{il} and $\bar{\gamma}_{il}$ are not both finite. In this case, we want that dimension’s proposal distribution G_{il} to have full support on the half or full real line, hence we set it to the exponential or double exponential. However, even if ϱ_{il} and $\bar{\gamma}_{il}$ are both finite, these outer bounds might still be very uninformative, and it will be undesirable to sample uniformly from such a large interval. If we don’t have $\chi_{il} < \varrho_{il} < \bar{\gamma}_{il} < \bar{\chi}_{il}$, then we still use the exponential or double exponential for G_{il} .

Sampling from the Conditional Distribution We take proposal draws from G_i conditional on \mathcal{G}_i using Gibbs sampling to draw one dimension at a time. Suppose we want to sample the s th new value for the l th cost parameter. Let $\gamma_{i,-l}^{(s)} = (\gamma_{i1}^{(s)}, \dots, \gamma_{i,l-1}^{(s)}, \gamma_{i,l+1}^{(s-1)}, \dots, \gamma_{i,n_\gamma}^{(s-1)})$ be the vector of current values for the other $L - 1$ dimensions. The set of values γ_{il} such that $(\gamma_{il}, \gamma_{i,-l}^{(s)}) \in \mathcal{G}_i$ is an interval, so we draw the new cost parameter $\gamma_{il}^{(s)}$ from G_{il} truncated to this interval.

We start by sampling 10,000 proposal draws, discarding the first half as burn-in and thinning the remainder by keeping every fifth draw. We end up with $S_\gamma^{\text{prop}} = 1,000$ proposal draws which are used to estimate the likelihood.

B.4.3 Metropolis-Hastings

Before running Metropolis-Hastings, we first estimate θ_γ via maximum (quasi-)likelihood. We start the maximum likelihood estimation at the following parameters: $\mu^{\text{volume}} = 0$, $\mu^{\text{density}} = 0$, $\mu^{\text{TU}} = (400, \dots, 400)$, $\mu^{\text{incumb}} = 0$, and $\Sigma = 50^2 I$. Let $\hat{\theta}_\gamma^{ML}$ denote the MLE point estimate, and let \hat{H}_γ^{ML} be the Hessian of the log-(quasi-)likelihood evaluated at $\hat{\theta}_\gamma^{ML}$. We use $\hat{\theta}_\gamma^{ML}$ as the starting point for Metropolis-Hastings.

For the jump distribution, we use a multivariate normal centered at zero, with variance-covariance matrix proportional to the inverse of the negative of the Hessian, $(-\hat{H}_\gamma^{ML})^{-1}$. The scaling parameter is chosen to target a rejection rate of between 0.4 and 0.6; in practice, we multiply the inverse Hessian by 0.03^2 . The inverse Hessian contains information about

how large a step to take in each dimension and how correlated the dimensions are. If the posterior marginal distribution of a parameter is very concentrated, then the corresponding diagonal element of the Hessian will be large, so the step size in that dimension will be small. When the posterior marginal distribution is very diffuse, the step size under the jump distribution will be large.

We start with 1 million Metropolis-Hastings iterations; then, we discard the first half as burn-in and thin the remainder by keeping every fiftieth draw. The result is $S_{\theta_\gamma} = 10,000$ draws of θ_γ . We report summary statistics for the sampled draws in tables [B.4](#) and [B.5](#).

Table B.4: Metropolis-Hastings Draws of Volume and Density Economy Parameters

	μ								σ			
	Small firms				Large firms							
	95% confidence set				95% confidence set				95% confidence set			
	Mean	SD	LB	UB	Mean	SD	LB	UB	Mean	SD	LB	UB
Volume 1	-11.507	21.928	-94.290	70.663	-9.275	26.542	-115.293	122.521	74.204	12.680	42.428	131.952
Volume 2	-30.158	11.590	-73.483	21.540	-8.649	14.055	-71.863	47.702	37.765	8.048	20.819	94.444
Volume 3	-19.683	11.886	-66.354	35.007	-19.900	11.456	-64.250	23.391	30.848	7.087	16.193	81.524
Volume 4	-9.298	37.740	-215.879	148.431	-25.790	18.891	-108.725	74.027	35.638	17.707	6.106	129.513
Density 1	-4.600	3.435	-18.132	8.625	2.711	4.137	-13.423	18.375	10.853	2.119	5.646	19.080
Density 2	0.178	6.126	-26.866	41.199	4.559	6.387	-25.001	34.046	14.120	4.128	5.934	40.203

Note: To construct the 95 percent confidence set, we compute the minimum and maximum values of each parameter among the 95 percent of Metropolis-Hastings draws (after burning and thinning) with the highest values of the likelihood.

Table B.5: Metropolis-Hastings Draws of TU Cost Parameters

	μ				σ			
	Mean	SD	95% confidence set		Mean	SD	95% confidence set	
			LB	UB			LB	UB
TU 401	462.882	6.223	434.242	492.351	25.258	5.491	11.646	59.539
TU 402	409.170	8.409	364.553	445.384	34.828	7.342	19.305	75.865
TU 403	500.153	4.549	480.217	519.402	18.029	4.019	8.191	44.645
TU 404	619.820	58.410	438.540	812.952	274.968	47.589	166.877	439.077
TU 405	549.370	14.984	488.505	623.228	62.880	11.855	33.318	126.407
TU 501	470.523	13.027	421.373	523.690	53.055	10.943	29.483	114.743
TU 502	371.022	14.678	303.596	416.804	60.811	10.796	32.745	104.638
TU 503	404.424	11.086	357.965	447.487	46.099	8.700	25.910	90.902
TU 504	389.384	8.013	348.499	423.183	31.509	7.748	15.528	93.869
TU 505	400.846	9.514	367.954	442.166	38.464	7.857	17.821	97.961
TU 506	411.759	7.021	374.353	435.900	25.834	10.239	11.028	82.689
TU 507	414.557	8.962	374.139	452.098	36.079	7.895	19.096	87.309
TU 508	368.979	6.619	344.987	395.647	23.611	5.007	11.442	46.705
TU 510	385.638	18.056	311.150	455.317	82.265	15.674	43.593	155.060
TU 511	395.946	8.124	369.086	429.932	31.944	6.946	13.101	76.150
TU 901	466.919	11.248	416.666	512.511	49.072	12.149	15.326	123.650
TU 902	480.806	13.166	427.306	539.300	53.675	12.224	27.554	139.168
TU 903	521.769	23.808	439.945	630.150	98.422	30.706	23.153	185.951
TU 904	416.109	5.616	394.391	448.530	22.078	4.863	11.771	52.692
TU 905	433.060	12.728	382.972	482.855	54.647	10.097	31.840	98.462
TU 906	453.025	8.670	396.626	490.295	31.768	10.166	15.857	102.798
TU 907	465.834	10.842	429.383	517.135	43.403	10.430	13.868	95.346
TU 908	516.200	19.083	400.057	591.895	73.687	16.649	40.145	156.052
TU 909	503.912	19.666	416.887	584.630	88.452	15.876	50.673	168.220
TU 1201	547.574	17.934	477.209	648.365	69.930	14.976	37.236	160.578
TU 1331	370.662	14.950	312.350	429.718	64.083	11.224	32.686	125.590
TU 1332	348.331	16.872	259.191	418.628	72.816	14.634	41.662	186.140
TU 1333	329.089	16.026	247.694	434.639	70.122	12.591	39.913	140.245
TU 1334	386.891	28.422	260.902	483.792	126.246	20.673	70.418	225.348

Table B.5: Metropolis-Hastings Draws of TU Cost Parameters

	μ				σ			
	Mean	SD	95% confidence set		Mean	SD	95% confidence set	
			LB	UB			LB	UB
TU 1335	369.221	11.345	327.185	409.970	47.260	8.444	24.269	89.210
TU 1336	365.385	8.698	326.605	406.966	34.658	6.933	17.774	70.070
TU 1339	399.456	14.771	339.603	465.194	63.102	11.609	34.540	121.559
Incumbency shifter	-9.196	8.353	-43.462	17.798				

Note: To construct the 95 percent confidence set, we compute the minimum and maximum values of each parameter among the 95 percent of Metropolis-Hastings draws (after burning and thinning) with the highest values of the likelihood.

B.4.4 Type Distribution Estimates


Across all firms and all TUs, the mean TU cost is 434 pesos per meal, compared to the mean TU base price of 464 pesos per meal discussed in section 5.3.1. The average markup from TU cost to TU base price is 7.7 percent of the TU base price. TUs on which firms bid higher base prices are also higher-cost to supply. Figure B.2 compares the two distributions separately for each TU.

B.5 Welfare Analysis

B.5.1 Sampling from the Conditional Type Distribution

For our welfare analysis, we need draws from the estimated type distribution $F_{\Gamma_i|z_i}(\cdot | \hat{\theta}_\gamma)$ conditional on type bounds \mathcal{G}_i for each firm i . Obtaining these draws requires sampling from a multivariate normal distribution truncated to a polyhedron. While it is possible to sample from truncated univariate normals in closed form using inverse transform sampling, there is no equivalent procedure in higher dimensions. Furthermore, naive rejection sampling is

Figure B.2: Distributions of Firms' TU Base Prices and Costs



Note: These violin plots show the distributions of firms' TU base prices $\tilde{\beta}_{ik}$ (left, in blue) and TU costs γ_{ik}^{TU} (right, in orange). In each TU k 's violin plots, an observation is a firm i . Firm i 's cost of supplying each TU k is computed from the mean of the estimated type distribution $F_{\Gamma_i|z_i}(\cdot | \hat{\theta}_\gamma)$ conditional on the type bounds \mathcal{G}_i .

computationally inefficient when the polyhedron’s volume is small relative to the dimension of the space: most of the sampled draws from the unconditional distribution are discarded, so the number of unconditional draws required in order to obtain each conditional draw is high.

Instead, we sample firm types γ_i via Gibbs sampling, drawing one dimension at a time as in appendix B.4.2. We run the Gibbs sampler for 100,000 iterations, discarding the first half as burn-in and thinning the remainder by a factor of five. We end up with $S_\gamma = 10,000$ draws of each firm’s cost parameters, which we use for the parametric results in Section 5.4.

B.5.2 Awarded Bid Cost and Markup Estimates

We present estimates of package costs and markups for each of the winning bids in table B.6.

Table B.6: Awarded Bid Cost and Markup Estimates

Firm	TUs	Meals (millions)	Price bid (pesos per meal)	Cost (pesos per meal)			Markup (% of price bid)		
				LB	UB	Mean	LB	UB	Mean
10	[401, 402, 901, 902, 903, 906, 908, 909]	21.17	462.98	415.80	462.98	450.56	0.00	10.19	2.68
13	[905, 1334]	5.23	386.07	270.69	386.07	330.80	-0.00	29.89	14.32
16	[504, 505, 507]	6.42	391.58	-582.71	391.58	329.92	0.00	248.81	15.74
17	[403, 404, 502, 503, 904, 907, 1201, 1331]	20.91	405.64	304.40	404.98	379.93	0.16	24.96	6.34
19	[1332, 1333, 1336]	9.48	340.05	338.56	339.90	339.49	0.04	0.44	0.17
26	[501]	1.64	459.07	441.15	459.07	453.69	0.00	3.90	1.17
28	[506, 508, 510, 511]	10.22	381.66	373.63	381.66	379.62	0.00	2.10	0.53
36	[1335, 1339]	3.60	381.77	323.21	381.77	370.04	-0.00	15.34	3.07
47	[405]	2.08	536.59	444.01	460.39	447.98	14.20	17.25	16.51